

# Nutrition and Health

## Comparison of total ionic strength adjustment buffers III and IV in the measurement of fluoride concentration of teas

Journal:	<i>Nutrition and Health</i>
Manuscript ID	NAH-17-0046.R1
Manuscript Type:	Original Manuscript
Date Submitted by the Author:	14-Nov-2017
Complete List of Authors:	Patel, Shilpa; Newcastle University, Dental school Omid, Narges; Newcastle University, Dental school Zohoori, Fatemeh; Teesside University, Maguire, Anne; Newcastle University Waldron, Kevin; Newcastle University, Institute for Cell and Molecular Biosciences Valentine, Ruth; Newcastle University, Dental school
Keywords:	Fluoride, Aluminium, TISAB, Caffein, Tea
Abstract:	<p>Background: Tea is the second most consumed drink in the UK and a primary source of hydration; it is an important source of dietary fluoride (F) for consumers and also abundant in aluminium (Al). Varying ranges of F concentrations in teas have been reported worldwide which may be, in part, due to differences in analytical techniques used to measure this ion. The effect of using total ionic adjustment buffers (TISAB) III or IV, when measuring F concentration of black teas available in the UK was investigated and compared. Based on this evaluation the effect of 3 different infusion times: 1 minute (1min), 10 min and 1hour (1h), caffeine content and tea form on the F contents of the tea samples was investigated.</p> <p>Methods: The F concentrations of 47 tea samples were measured directly using a fluoride ion-selective electrode, TISAB III and IV, and infusion times of 1min, 10min and 1h.</p> <p>Results: Mean (SD) F concentration of tea samples for all infusion times was statistically significantly higher (<math>p &lt; 0.001</math>) measured by TISAB IV (4.37 (2.16) mg/l) compared with TISAB III (3.54 (1.65) mg/l). A statistically significant positive correlation (<math>P &lt; 0.001</math>) was found between Al concentration (mg/l) and differences in F concentration (mg/l) measured using the two TISABs; the difference in F concentration measured by the two TISABs increased with the magnitude of Al concentration.</p> <p>Conclusion: Due to higher concentrations of F and Al in teas and their complexing potential, use of TISAB IV facilitates more accurate measurement of F concentration when using F-ISE and a direct method.</p>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

SCHOLARONE™  
Manuscripts

For Peer Review

1  
2  
3 **Comparison of total ionic strength adjustment buffers III and IV in the measurement of**  
4 **fluoride concentration of teas**  
5  
6

7 Shilpa Patel<sup>1</sup>, Narges Omid<sup>1</sup>, Fatemeh V Zohoori<sup>2</sup>, Anne Maguire<sup>1</sup>, Kevin J Waldron<sup>3</sup> and  
8 Ruth A Valentine <sup>\*1,3</sup>  
9  
10

11  
12  
13  
14 <sup>1</sup> Centre for Oral Health Research, School of Dental Sciences, Newcastle University, UK  
15

16 <sup>2</sup> School of Health and Social Care, Teesside University, UK  
17

18 <sup>3</sup> Institute for Cell and Molecular Biosciences, Newcastle University, UK  
19  
20

21 **Running title:** Comparison of two buffers for measuring fluoride concentration of teas  
22

23  
24 \* Corresponding author:  
25

26 Ruth A Valentine  
27

28  
29 School of Dental Sciences, Newcastle University, NE2 4BW, UK  
30

31 Email: [Ruth.valentine@ncl.ac.uk](mailto:Ruth.valentine@ncl.ac.uk)  
32  
33

34  
35  
36 **Acknowledgment**  
37

38 The present study was partially supported by a grant from the Borrow Foundation. K.J.W.  
39 was supported by a Sir Henry Dale Fellowship funded by the Wellcome Trust and the Royal  
40 Society (098375/Z/12/Z).  
41  
42

43 The views expressed in this paper are those of the authors and not those of the funding body.  
44

45 RV, AM and VZ designed the study. SP collected and analysed samples. KW analysed AI  
46 samples. NO, was involved in re-analysis. RV, AM, VZ, NO and KW were involved in  
47 manuscript drafting. All authors read and approved the final manuscript.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Abstract

Background: Tea is the second most consumed drink in the UK and a primary source of hydration; it is an important source of dietary fluoride (F) for consumers and also abundant in aluminium (Al). Varying ranges of F concentrations in teas have been reported worldwide which may be, in part, due to differences in analytical techniques used to measure this ion.

The effect of using total ionic adjustment buffers (TISAB) III or IV, when measuring F concentration of black teas available in the UK was investigated and compared. Based on this evaluation the effect of 3 different infusion times: 1 min, 10 min and 1 h, caffeine content and tea form on the F contents of the tea samples was investigated.

Methods: The F concentrations of 47 tea samples were measured directly using a fluoride ion-selective electrode, TISAB III and IV, and infusion times of 1min, 10min and 1h.

Results: Mean (SD) F concentration of tea samples for all infusion times was statistically significantly higher ( $p < 0.001$ ) measured by TISAB IV (4.37 (2.16) mg/l) compared with TISAB III (3.54 (1.65) mg/l). A statistically significant positive correlation ( $P < 0.001$ ) was found between Al concentration (mg/l) and differences in F concentration (mg/l) measured using the two TISABs; the difference in F concentration measured by the two TISABs increased with the magnitude of Al concentration.

1  
2  
3  
4  
5  
6  
7  
8  
9 Conclusion: Due to higher concentrations of F and Al in teas and their complexing  
10 potential, use of TISAB IV facilitates more accurate measurement of F concentration  
11 when using F-ISE and a direct method.  
12  
13  
14  
15

16  
17 **Keywords:**

18  
19  
20 Tea, fluoride, aluminium, TISAB, caffeine  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Introduction

The tea plant is known to have a high bio-accumulation of both fluoride (F) and aluminium (Al), absorbing high levels from acidic soils at high temperatures. Both elements are accumulated mainly in leaves, with higher concentrations found in mature leaves. F accumulation has been reported to range from 300-1000 mg/kg in young leaves and from 600-2800 mg/kg in old leaves (Fung et al., 1999) while Al ion concentrations from 997 to 5600 mg/kg have been found in young and old leaves respectively (Wong et al., 1998). Complexes of F with Al occur in acidic soils (pH<5.5) and form  $\text{AlF}^{2+}$ ,  $\text{AlF}_2^+$  or  $\text{AlF}_3$ , with the latter being the dominant form (Fung et al., 1999).

In contrast to the concentrations found in tea leaves themselves, the amounts of F and Al ions released to tea liquors during infusion are much lower. For example, for dry tea products containing 300-2677 mg/kg Al and 107-878 mg/kg F, the concentrations in tea infusions were reported at 0.7-3.5 mg/l (Wong et al., 1998) and 0.6-1.9 mg/l (Fung et al., 1999).

Optimal exposure to F has benefits for oral health through prevention of dental caries and encouragement of bone and teeth remineralisation (McArthur, 2014; World Health Organization, 1994; Fung et al., 1999). These oral health benefits are best achieved through daily consumption of drinking waters with low F concentrations of 0.8 – 1.0 mg/l (World Health Organization, 2010). However, chronic excessive systemic (body)

1  
2  
3  
4  
5  
6  
7  
8  
9 exposure to F can have health implications, including increased risk of dental fluorosis  
10 (mottled spots seen on tooth enamel) if the excessive exposure is during the “window of  
11 susceptibility” of early childhood, and skeletal fluorosis, a crippling condition leading to  
12 an increase in bone mass if the excessive exposure extends into and through adulthood  
13 (World Health Organization, 1994; Medical Research Council, 2002; Den Besten and  
14 Li, 2011; Joshi et al., 2011). **An upper tolerable intake of 10mg of F per day for adults  
15 aged 19 years or above has been recommended with a Dietary Reference Intake (DRI)  
16 of 4 mg/day (Otten et al., 2006)**

17  
18  
19  
20  
21  
22  
23  
24  
25  
26 Exposure to Al is limited but can occur from drinking waters, in which it is used as a  
27 coagulant to reduce organic matter, as well as from foods, usually due to Al-containing  
28 food additives and from antacids. Al has been suggested as a possible risk factor for  
29 dementia promotion, onset and progression (Dolara, 2014; Henderson et al., 2002;  
30 Flaten, 2002), but the evidence remains equivocal. **The Joint Food and Agriculture  
31 Organization (FAO) / World Health Organization (WHO) Expert Committee on Food  
32 Additives and Food Contaminants have recommended a provisional tolerable weekly  
33 intake (PTWI) of Al of 1.0 mg/kg bw. (FAO/WHO, 2006).** The bioavailabilities of F  
34 and Al are dependent upon their chemical forms. Regarding F, its bioavailabilities in  
35 water, tea and solid foods differ; being up to 100% in water, 94.9% in teas (World  
36 Health Organisation, 1984) and between 50-80% in solid foods, depending on their  
37 chemical composition (Grooper et al., 2009). The same trend is seen for aluminium,  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9 with systemic uptake of Al being generally very low, but higher from water (0.28% of  
10 intake absorbed) and from teas (0.37%) than from solid foods (0.1% of intake absorbed)  
11 (Dolara, 2014) (Yokel and Florence, 2008). Since tea is the second most consumed  
12 drink worldwide after water, among both adults and children (Emekli-Alturfan et al.,  
13 2009), it may form a significant contribution to an individual's F and Al intake. In view  
14 of this, many studies have measured F and Al contents of different types of teas  
15 worldwide.  
16  
17

18  
19 Globally, black tea infusions have been reported to vary greatly in F content, ranging  
20 from 0.57 – 6.01 mg/l and in Al content, from 0.065 – 15 mg/l (Flaten, 2002; Yi and  
21 Cao, 2008; Wong et al., 2003; Emekli-Alturfan et al., 2009; Cao et al., 2006; Chan et  
22 al., 2013). These variations primarily reflect the local geological, cultivation and  
23 processing conditions as well as the different forms of teas and brewing times used.  
24 However, in addition, the varying ranges of F and Al concentrations reported may be, in  
25 part, due to differences in the analytical techniques used. For F these include ion  
26 chromatography (IC), spectrophotometry and fluoride ion selective electrode (F-ISE)  
27 with the latter being used in the majority of studies. In contrast, for Al measurement the  
28 methods used range from graphite furnace atomic absorption spectrometry (GFAAS)  
29 and inductively coupled plasma atomic emission spectrometry (ICP-AES) to inductively  
30 coupled plasma mass spectrometry (ICP-MS), with ICP-AES being the primary method  
31 used (Flaten, 2002).  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

However, since the F-ISE responds only to free F ions, samples which contain free or ionisable F ions should be analysed directly after adding total ionic adjustment buffer (TISAB) (Martínez-Mier et al., 2011). This buffer has an important role in reducing interference from certain elements (e.g. aluminium, iron and silicon). The purpose of adding TISAB to samples is threefold; i) and ii) to adjust the sample and standards to the same ionic strength and pH, and; iii) to de-complex F ions from certain interfering elements. TISABs are commonly used in their commercially available forms, namely II, III and IV. TISAB II and III adjust the pH of samples and standards to between 5 and 5.5, while TISAB IV adjusts to a pH of approximately 8.5. All TISABs contain an Al ion masking agent; cyclo-hexylene-dinitrilotetraacetic acid (CDTA), which complexes with Al preferentially, freeing F from complexes with this ion (Schamschula et al., 1985; Tušl, 1970). TISAB II and III can complex approximately 5 mg/l of Al in a sample containing 1 mg/l of F, and the ratios used to buffer the samples are 1:1 and 1:10 for TISAB II and III respectively, while TISAB IV is recommended for use in samples with higher concentrations of Al ion since it contains stronger chelating agents and eliminates interference from Al-F complexes (Thermo Scientific, Orion, User Guide 2014; Caslab 1996). According to the literature, the most commonly used method for F analysis of teas is a direct method (Cao et al., 2006; Chan et al., 2013; Emekli-Alturfan et al., 2009; Fung et al., 1999; Koblar et al., 2012; Malinowska et al., 2008). However these studies have used TISAB II (Chan et al., 2013) or III (Emekli-Alturfan et al.,

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

2009), or have not reported the type of TISAB used (Cao et al., 2006; 2009; Fung et al., 1999; Koblar et al., 2012; Malinowska et al., 2008). In view of the high Al concentrations of up to 13 mg/l reported for tea samples (Koch et al., 1988; Erdemoglu and Gucer, 2005), measurements of F in teas reported in the literature might be underestimated due to types of TISABs used, which do not allow for the high Al content of some teas.

These high concentrations of both F and Al in teas, as well as the wide range of F concentrations in teas reported in the literature, highlight the importance of using standardised measurement procedures when using a F-ISE and a direct method. Therefore the main aim of this study was to investigate and compare the effect of using two different TISABs; III and IV, when measuring F concentration in black teas, commercially available in the UK, taking into account the Al concentrations. Based on the preferred/appropriate TISAB, the subsidiary aim was to investigate the effect of 3 different infusion times; 1 min, 10 min and 1 h, as well as the caffeine content and form of tea on the F concentrations of the tea samples

## Method

### *Sourcing samples*

In total, 47 black teas were identified through a thorough search of supermarket websites and/or visiting the supermarkets directly. The teas were purchased from the following UK supermarkets; Asda, Morrisons, Sainsbury's and Tesco in March 2014

1  
2  
3  
4  
5  
6  
7  
8  
9 and samples categorised based on: (i) caffeine content; caffeinated or decaffeinated, and  
10  
11 ii) type of packaging; teabag, loose (tea leaves) and instant tea.  
12

### 13 *Sample Preparation*

14  
15 Tea samples were prepared to represent infusions ready for consumption, simulating the  
16  
17 customary habits of tea drinkers (Ruxton and Hart, 2011). **A longer non-representative**  
18  
19 **time point of 1h was also included to look at a maximal effect of infusion on F content.**  
20  
21

22 For tea bags and loose teas, where the infusion times could be measured, infusion times  
23  
24 of 1 min, 10 min and 1 h were used.  
25

26 Tea Bags: 3 whole tea bags (one for each time point) were separately infused with 200  
27  
28 ml of boiling double de-ionised water (DDiH<sub>2</sub>O) for 1 min, 10 min and 1 h; tea bags  
29  
30 were not squeezed or stirred.  
31

32  
33 Loose Tea: Manufacturers' recommendations of 1 level teaspoon (2.20 g) of dry leaves  
34  
35 per cup were followed and 200 ml boiling DDiH<sub>2</sub>O added prior to 3 separate infusions  
36  
37 timed at 1 min, 10 min and 1 h. The infusions were then strained with a stainless steel  
38  
39 tea strainer, removing loose leaves to prevent further infusion and the strainer rinsed  
40  
41 with DDiH<sub>2</sub>O twice between each use.  
42  
43

44 Instant Tea: Product recommendations of 1 teaspoon per cup (1.00 g) were used. Due to  
45  
46 the tea form, an infusion time was not applicable, therefore one infusion was prepared  
47  
48 per sample. One gram of tea granules was weighed and 200 ml boiling DDiH<sub>2</sub>O added.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9 All prepared tea liquors were cooled to room temperature, stirred and transferred to 30  
10 ml polystyrene universal containers. For F analysis the infusions were freshly prepared  
11 on a daily basis and analysed on the same day. For Al analysis, samples were mixed  
12 thoroughly and frozen prior to analysis.  
13  
14  
15  
16

#### 17 *F analysis of tea liquors*

18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
Tea samples were analysed for F by a direct method (Martínez-Mier et al., 2011) and in  
triplicate using a F-ISE (Model 9409; Orion) and meter (Model 900A; Orion) after  
sample buffering with either TISAB III (Orion 940911 Thermo Scientific USA) or  
TISAB IV (Fluka analytical, Sigma-Aldrich, Germany). The validity of the method was  
determined by adding a known amount of F to 10% of both the TISAB III and TISAB  
IV buffered samples analysed and measuring the F recovery. In addition, 10% of all tea  
samples were re-analysed on a different day to check the reliability of the method.

#### 59 *Determination of Al Content*

60  
All frozen tea liquor samples were thawed to room temperature before analysis. ICP-  
MS was used to determine the Al content of the tea liquors using a Thermo X-series  
instrument operating in standard mode. For each analysis, 500 µl of tea sample was  
diluted 10-fold with 2.5% (w/v) high-purity HNO<sub>3</sub> (Merck) containing 20 µg/l silver  
(Ag) and platinum (Pt) as internal elemental standards. For each sample, aluminium  
(<sup>27</sup>Al), silver (<sup>107</sup>Ag) and platinum (<sup>195</sup>Pt) isotopes were monitored sequentially using  
the peak-jump method (100 individual reads of 20 milliseconds on each isotope, across

1  
2  
3  
4  
5  
6  
7  
8  
9 3 channels of 0.02 atomic mass unit [AMU] separation, each in triplicate). Al  
10 concentrations were determined by comparison with matrix-matched standard solutions  
11 of known elemental composition (0, 1, 5, 10, 50, 100, 500, 1000 and 5000 mg/l)  
12 analysed within the same run.  
13  
14

#### 15 16 17 *Data handling and statistical analysis* 18

19 To allow comparison of the performance of TISAB III and IV at lower and higher Al  
20 concentrations, F data were grouped according to the Al concentration of the samples;  
21  $\leq 5$  mg/l and  $>5$  mg/l. Data were analysed using SPSS version 22 (SPSS, Chicago, IL,  
22 USA). Descriptive analysis was carried out to report mean (SD) and median (maximum  
23 and minimum) values to allow comparison with other studies. Data were treated  
24 according to the results of their normality testing. A Paired t-test was performed to  
25 compare all F concentrations measured by TISAB III and IV (n=137). **Two way**  
26 **ANOVA with Sidak's multiple comparison test was used to compare F concentration of**  
27 **tea samples between the two types of TISAB used.** Regression analysis was used to  
28 investigate the effect of Al concentration on the difference between measured F  
29 concentration using TISAB III and IV. Mann Whitney U-tests were used to investigate  
30 the effect of caffeine content and tea form on the F concentration of tea samples and a  
31 Kruskal-Wallis test, followed by pairwise multiple comparison analysis was performed  
32 to investigate the effect of infusion time on F concentration.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Results

The mean (SD) recovery of F added to a randomly selected sub group of 10% of tea samples was 98.5 (3.1)% and 98.0 (2.9) % using TISAB III and TISAB IV respectively, indicating satisfactory validity of the F analysis method used. In addition, the reliability of the method was checked by re-analysis of 10% of tea samples using both TISAB III and IV, which showed <4% difference between analysis and immediate re-analysis. A small number of tea samples were analysed immediately after infusion and again after 24 hours and 48 hours of storage at 4°C. Samples were incubated in the fridge and analysed at room temperature. Analysis with TISAB III resulted in a significant decrease in F concentration 48 hours post preparation. Re-analysis with TISAB IV after 48 hours storage showed no significant change in F concentration with time (results not shown). The reduction in F concentration measured with TISAB III is most probably due to the phenomenon of 'tea creaming' (Jobstl et al., 2005), a naturally occurring precipitation effect that occurs during cooling due to some of the components that are soluble in hot water, being insoluble in cold water and precipitating (approximately 30% of the total solids). Due to the possible 'creaming' effect after sample storage at 4°C, teas were analysed immediately after tea liquor sample preparation. As Table 1 shows, mean (SD) F concentration (mg/l) for 45 teas measured at all 3 infusion times and the 2 instant tea samples measured at one infusion time (n=137) by TISAB III and IV were 3.54 (1.65) and 4.37 (2.16) mg/l respectively. A

1  
2  
3  
4  
5  
6  
7  
8  
9 statistically significantly ( $P < 0.001$ ) higher F concentration was measured when TISAB  
10 IV was used.

11  
12 A statistically significant positive correlation ( $R^2 = 0.2666$ ,  $P < 0.001$ ) was found between  
13 Al concentration (mg/l) and the differences between F concentration (mg/l) measured  
14 using the two TISABs (Figure 1) with “between-method differences in measured F  
15 concentration (mg/l) =  $0.123 \times \text{Al concentration (mg/l)}$ ”. This highly significant  
16 correlation suggests that the difference in F concentration measured by the two TISABs  
17 increased with the magnitude of Al concentration: i.e. an increase in the difference in F  
18 concentration measured between the two TISABs of 0.123 mg/l for every 1 unit  
19 increase in Al concentration (mg/l).  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29

30  
31 Figure 2 compares the mean measured F concentration (mg/l) using TISAB III and IV  
32 according to the Al concentration ( $< 5$  mg/l and  $> 5$  mg/l) of the samples. For tea samples  
33 with a Al concentration of  $\leq 5$  mg/l, the difference in mean F concentration (mg/l)  
34 measured using TISAB III and IV was low (mean (SD) F concentration: 2.30 (0.86)  
35 mg/l and 2.44 (0.89) mg/l, respectively; ns), whereas for tea samples with a Al  
36 concentration of  $> 5$  mg/l, the corresponding difference was relatively high (mean (SD)  
37 F concentration: 4.16 (1.62) mg/l using TISAB III and 5.33 (1.98) mg/l using TISAB  
38 IV;  $P < 0.001$ ).  
39  
40  
41  
42  
43  
44  
45  
46  
47

48 Since the measured F concentrations were statistically significantly higher in samples  
49 analysed using the TISAB IV buffer compared with TISAB III, the TISAB IV was our  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9 preferred buffer for the F analysis of tea and therefore only analyses and comparisons of  
10 teas for TISAB IV based samples are presented in the following sections.

11  
12 Information on the brand/names, form of teas (e.g. loose, tea bags and instant), caffeine  
13 content and F concentration of all tea samples at infusion times of 1 min, 10 min and 1 h  
14 is presented in Table 2. The majority of the 47 analysed tea samples were caffeinated  
15 (n=39) and in the form of teabags (n=37). The F concentrations of the 2 instant teas  
16 were 7.78 and 9.91 mg/l for PG Tips and Tetley pure tea granules respectively. Among  
17 the remaining samples, the highest F concentration (9.44 mg/l) was found for PG Tips  
18 (de-caffeinated) after 1 h infusion, and the lowest (1.06 mg/l) for Twinings Assam  
19 (caffeinated) after 1 min of infusion.

20  
21 Based on infusion time, the mean (SD) F concentration of tea samples infused for 1 h  
22 was higher [5.17 (2.27) mg/l] than those infused at 10 min [4.52 (2.11) mg/l] and 1 min  
23 [3.16 (1.36) mg/l], and the difference between infusions at 1 min and 10 min as well as  
24 those infused at 1 min and 1 h was statistically significant ( $p < 0.001$ ) (Table 3). Data for  
25 instant teas are presented separately since they could only be prepared at a single time  
26 point.

27  
28 Based on caffeine content, across the 3 infusion times the F concentrations of de-  
29 caffeinated teas were significantly higher ( $p < 0.05$ ) than caffeinated teas (Table 4). On  
30 the same basis, across all infusion times and as Table 4 also describes, a significantly  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

( $p < 0.001$ ) higher F concentration was found in teabags (4.62 (1.86) mg/l) compared with loose teas (2.82 (0.87) mg/l).

### **Discussion and conclusion**

This study measured F concentration of a range of teas available in the UK market using a direct method and F-ISE. The main aim was to investigate and compare the effect of using two different TISABs; III and IV.

According to the literature, the preferred method for ionic F analysis of teas is a direct method using F-ISE (Cao et al., 2006; Chan et al., 2013; Emekli-Alturfan et al., 2009; Fung et al., 1999; Koblar et al., 2012; Malinowska et al., 2008; Chandrajith et al., 2007). Although an overnight acid-diffusion pre-treatment can be used for samples containing non-ionic F complexes, which releases F ions from such complexes prior to measurement using F-ISE (Taves, 1968; Soto-Rojas et al., 2005), the use of a direct method was preferred. This is mainly because >90% of F in tea infusions exists in the ionic form which can be measured directly after adding a TISAB buffer (World Health Organisation, 1984; Fung et al., 1999). The majority of studies have used a TISAB:sample ratio of 1:1, and most studies have primarily used TISAB II. However, the reported high levels of Al in tea samples, and the complexing interaction of Al with F, suggest that these studies may have underestimated F concentrations in view of the specific TISABs used. The statistically significantly higher concentration of F

1  
2  
3  
4  
5  
6  
7  
8  
9 measured when using TISAB IV compared with TISAB III shows that TISAB IV is  
10 more capable of releasing F ions due to the reduction in the stability of the Al-F  
11 complex through adjustment of the sample pH to 8.5.  
12

13  
14  
15 The effect of TISAB IV in releasing a greater proportion of the F ions was further  
16 confirmed with the statistically significant positive correlation ( $P < 0.001$ ) found between  
17 Al concentration (mg/l) and difference in F concentration (mg/l) measured using the two  
18 TISABs. The study found an increase in the difference in F concentration of  $>0.1$  mg/l  
19 for every 1 unit increase in Al concentration (mg/l). The manufacturer's recommended  
20 limits of measurement when using TISAB III at 1 ppm F concentration are 5 mg/l Al  
21 (Thermo Scientific Orion ion plus Fluoride Electrode, Instruction Manual, 2014).  
22 Interestingly, in these tea samples, all but one contained more than 1 ppm F (1 mg/l),  
23 indicating that TISAB III may not be a suitable buffer to measure F in tea infusions,  
24 even with relatively low Al concentrations.  
25  
26

27  
28  
29 Previous reports have determined F concentrations of black tea infusions ranging from  
30 low levels identified in Brazilian tea samples (0.13 mg/l) (Hayacibara et al., 2004) to  
31 high levels measured in UK sourced teas (8.85 mg/l); both studies used ISE with TISAB  
32 II (Chan et al., 2013). The F concentrations found in teas purchased in the UK for this  
33 present study fit within and extend beyond this range, but the results obtained are in  
34 agreement with previous outcomes; that F does accumulate in the tea leaves and that its  
35 content in tea liquors increases with the duration of infusion (Chan et al., 2013; Quock  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9 et al., 2012; Koblar et al., 2012). However, the statistically significant positive  
10 correlation observed between F and Al concentrations of the teas analysed highlights  
11 the importance of using the correct buffer prior to measurement. Although F in tea has  
12 been extensively investigated and analysed, the range of concentrations previously  
13 proposed may be lower than the actual total available F concentrations (i.e. the ionic F  
14 content available for absorption in the gastro-intestinal tract (GIT)) present in black  
15 teas. Concentrations of F in tea are dependent upon the original source of the tea plant  
16 as a raw material, its maturity and the pH of the soil in which it has grown (Janiszewska  
17 and Balcerzak, 2013). Many popular brands and supermarket varieties of teas are  
18 produced through mixing of *Camellia sinensis L.* blends derived from various  
19 geographical locations (Hayacibara et al., 2004) (Table 2). Pure blends from Assam and  
20 Ceylon had the lowest F concentration in comparison with all other teas, ranging from  
21 0.94 – 1.32 mg/l for tea liquors. These data are also in line with studies analysing  
22 similar pure blend teas in the UK and Europe, prepared with shorter infusion times  
23 (Chan et al., 2013; Chandrajith et al., 2007) as well as the upper limit ranges found in  
24 Ceylon tea analysis (0.32 – 1.68 mg/l respectively) (Chandrajith et al., 2007). Reports in  
25 the UK and China have suggested that economy blend teas and teas with no claim of tea  
26 plant country of origin have the highest concentrations of F (Cao et al., 2006), and  
27 previous studies focusing on economy teas in the UK support this, recording the F  
28 concentration of economy ‘value’ teas to range from 6.20 - 8.85 mg/l (Sainsbury Basics  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9 – Tesco Value respectively) when infused for up to 30min (Chan et al., 2013). These  
10 results are in agreement with the current study in which the F content of economy teas  
11 ranged from 2.88 – 9.00 mg/l at 1 h infusion, and were some of the highest F  
12 concentrations recorded. Economy branded teas are manufactured with older leaves,  
13 branches and roots of the tea plant (Wong et al., 2003); these mature parts of the tea  
14 plant are estimated to contain up to 10-fold more F than the younger leaves and buds  
15 (Fung et al., 1999).  
16

17  
18 Previous studies investigating Al content independently, as well as investigations of F  
19 and Al contents together, have found the Al content in black teas to range from 0.065  
20 (Rao and Rao, 1994) to 13 mg/l (Erdemoglu and Gucer, 2005; Koch et al., 1988). These  
21 studies applied different measurement methods including atomic emission spectrometry  
22 (AES), inductively coupled plasma atomic emission spectrometry (ICP-AES), flame  
23 atomic absorption spectrometer (FAAS), graphite furnace atomic absorption  
24 spectrometry (GFAAS) and ICP-MS. In the present study ICP-MS was used and found  
25 the Al concentration for the combined infusion times (n=137) ranged from 1.7 to 21.9  
26 mg/l which is wider than the 1-6 mg/l previously reported by other studies (Flaten,  
27 2002). This may partly be explained by differences in the analytical methods used, for  
28 example, a previous study investigating Al content by ICP-AES of Tetley black tea  
29 infused for 3 min in 250 ml boiling deionised water found 2.90 mg/l (Powell et al.,  
30 1993) while the current study determined Al concentration at 7.46 mg/l in Tetley  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9 Original tea bags infused for 1min. Another possible explanation for this variability  
10 may be the year to year variation in multiple crops used to blend this tea. 'Tea  
11 creaming' may also influence levels of Al measured in teas, although the addition of  
12 HNO<sub>3</sub> to the sample in a digestion step prior to ICP-MS analysis should have allowed  
13 Al to be released from any insoluble material and measured. The higher concentrations  
14 determined in this study may be indicative of more sensitive detection limits of the ICP-  
15 MS in comparison with methods used in previous studies, or may be explained in some  
16 part by the variation between different batches used to blend the tea product.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

27 In terms of the effect of infusion time on F content of teas, the current study used  
28 different infusion times of 1 min, 10 min and 1 h to cover the range of infusion times  
29 previously measured by other studies (up to 10 min), as well as to mirror customary  
30 home tea preparation (shortened infusion time of 1 min). A longer non-representative  
31 time point of 1 h was included to look at a maximal effect of infusion of F content.  
32  
33  
34  
35  
36  
37

38 A wide range of infusion times, between 2 min (Chan et al., 2013; Wong et al., 2003)  
39 and 6 h (Wong et al., 2003; Fung et al., 1999), has been reported in the literature for the  
40 preparation of tea infusions, with the resultant mean F concentrations ranging from 0.71  
41 mg/l (Chan et al., 2013) to 1.89 mg/l (Fung et al., 1999) for samples prepared at 2 min  
42 and 6 h respectively. The widest range in F concentrations was reported by Chan et al.  
43 between 0.71 and 8.80 mg /l for samples infused for 2 min and 30 min, respectively  
44 (Chan et al., 2013). However, another study, despite using a wider infusion time range  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9 (up to 6 h), reported a narrower range of F concentrations, between 0.91 mg/l at 5 min  
10 and 1.90 mg/l at 2 h time points (Fung et al., 1999). That range was smaller than the  
11 1.06 mg/l to 9.44 mg/l range observed in this present study for samples infused at 1 min  
12 and 1 h, respectively, which are more in line with the range reported by Chan et al  
13 (Chan et al., 2013) in which a significantly higher F concentration was recorded after 30  
14 min infusion.  
15  
16

17  
18  
19  
20  
21 It has been reported that during the first 30 min of infusion a large amount of F is  
22 released into the tea liquor, with 77-85% being released in the first 5 min and the  
23 amount of released F is directly proportional to the duration of infusion (Fung et al.,  
24 1999). The findings of the present study show that 61% of F was released in the first 1  
25 min of infusion, and 88% in the first 10 min, with the amount of measured F statistically  
26 significantly increasing between 1 min and 10 min and between 1 min and 1 h, while  
27 the difference between 10 min and 1 h was not statistically significant.  
28  
29

30  
31  
32  
33  
34  
35  
36  
37 In terms of the caffeine content of the teas, statistically significantly higher F  
38 concentrations were found in decaffeinated brands compared with caffeinated varieties,  
39 a finding previously reported along with one possible explanation; the use of fluoridated  
40 water in the decaffeinating process (Yi and Cao, 2008; Chan and Koh, 1996). This  
41 process involves removing caffeine from the tea by soaking the tea leaves in hot water  
42 for a period of time, filtering the solution through carbon to remove caffeine and  
43 returning the water to the tea for reabsorption of flavours and oils, however this is not  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8 generally used commercially. Another possible explanation is the known effect of  
9 caffeine on tea creaming. Tea with lower levels of caffeine produce less 'creaming'  
10 (Jobstl et al., 2005), and therefore potentially a lower loss of minerals during the cooling  
11 process, which may account for some of the variation observed in this study.  
12  
13  
14  
15  
16

17 Fermentation is a vital stage in black tea manufacture, and is also characteristic of teas  
18 with elevated F concentrations (Janiszewska and Balcerzak, 2013; Malinowska et al.,  
19 2008). It has been suggested that this may be due, in part, to greater amounts of  
20 polyphenols (enhancing fermentation) present in the tea plant, however it has also been  
21 shown that polyphenols correlate negatively with F concentrations, with older plants  
22 having lower polyphenol but higher F concentrations (Lu et al., 2004). With regard to  
23 instant teas, freeze dry processing involves concentrating the extract from brewed pure  
24 tea leaves to produce a powdered form. The highly concentrated nature of the tea extract  
25 itself is the probable reason for the highest F concentrations being determined in the  
26 instant tea infusions when compared with loose tea and tea bags at 2.66 mg/l and 4.65  
27 mg/l respectively.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 Regarding diet, the systemic bioavailability of F is considered to be high, between 50-  
42 80% in the fed state and 100% in a fasted state (Ophaug, 1990). F presents a biphasic  
43 nature in the form of benefits and risks to dental and general health and as a source of  
44 topical F in the mouth and systemic F when absorbed from the GIT (Simpson et al.,  
45 2001). Tea, together with dentifrices such as toothpaste and fluoridated water, could  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9 have both positive and negative health benefits depending on the overall level of  
10 consumption. Once absorbed systemically, 90% of the F retained in the body is  
11 incorporated into calcified tissues including bone and enamel crystallites (World Health  
12 Organization, 1994). The presence of Al reduces F bioavailability, while a deficiency in  
13 Al may heighten F uptake from the GIT (Den Besten and Li, 2011). The Al content of  
14 black teas, together with the positive correlation between F and Al concentrations, may  
15 reduce the body's ability to absorb and utilise the F that tea infusions contain. In terms  
16 of F, a Dietary Reference Intake (DRI) of 4 mg/day and upper tolerable intake of 10  
17 mg/day for adults aged 19 years and above has been recommended (Otten et al., 2006)  
18 and according to the UK National Diet and Nutrition Survey (Henderson et al., 2002)  
19 the average daily consumption of tea in the UK is 540 ml (2-3 servings). This indicates  
20 that, based upon the F content of teas found in the present study, levels of daily F  
21 consumption in a UK population living in optimally fluoridated areas could be towards  
22 the upper tolerable range. The current *in vitro* study used de-ionised water to prepare  
23 infusions, in contrast to the situation found *in vivo* where higher levels of F may be  
24 consumed in those areas where fluoridated water is available and used for tea making  
25 (Koblar et al., 2012).  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

46 Since teas may comprise a significant proportion of individual's daily fluid intake, it is  
47 important to increase public awareness on any health implications of drinking excessive  
48 amounts of tea, when informing on the need for balance between benefit and risk. In  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7  
8  
9 addition, health professionals should take tea consumption among young children into  
10 consideration prior to prescribing F supplements. It is noted by the authors, however,  
11 that there is no evidence in Westernised society that consuming commercially available  
12 black tea increases the risk of developing fluorosis.  
13  
14  
15  
16

17  
18 Due to the high concentration of F and Al in the tea samples analysed and their potential  
19 for complexing with F, the use of TISAB IV provides a more accurate measure of F  
20 concentration when using a direct method and F-ISE.  
21  
22  
23

24  
25 The wide range of F concentrations found in popular teas available in the UK together  
26 with fluoridated tap water being available for the preparation of tea infusions in some  
27 areas, highlights the need for further information on the typical amounts of F consumed  
28 through this beverage.  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**References:**

- Cao J, Zhao Y, Li Y, et al. (2006) Fluoride levels in various black tea commodities: Measurement and safety evaluation. *Food and Chemical Toxicology* 44: 1131-1137.
- Chan JT and Koh SH. (1996) Fluoride content in caffeinated, decaffeinated and herbal teas. *Caries Research* 30: 88-92.
- Chan L, Mehra A, Saikat S, et al. (2013) Human exposure assessment of fluoride from tea (*Camellia sinensis* L.): A UK based issue? *Food Research International* 51: 564-570.
- Chandrajith R, Abeyapala U, Dissanayake CB, et al. (2007) Fluoride in Ceylon tea and its implications to dental health. *Environmental geochemistry and health* 29: 429-434.
- Den Besten PK and Li W. (2011) Chronic Fluoride Toxicity: dental Fluorosis. In: Buzalaf MA (ed) *Fluoride Intake and the Oral Environment* Basel, Switzerland: Karger, 81-96.
- Dolara P. (2014) Occurrence, exposure, effects, recommended intake and possible dietary use of selected trace compounds (aluminium, bismuth, cobalt, gold, lithium, nickel, silver). *International Journal of Food Sciences and Nutrition* 65: 911-924.
- Emekli-Alturfan E, Yarat A and Akyuz S. (2009) Fluoride levels in various black tea, herbal and fruit infusions consumed in Turkey. *Food and Chemical Toxicology* 47: 1495-1498.
- Erdemoglu SB and Gucer S. (2005) Selective determination of aluminum bound with tannin in tea infusion. *Analytical Sciences* 21: 1005-1008.
- Flaten TP. (2002) Aluminium in tea—concentrations, speciation and bioavailability. *Coordination Chemistry Reviews* 228: 385-395.
- Fung KF, Zhang ZQ, Wong JWC, et al. (1999) Fluoride contents in tea and soil from tea plantations and the release of fluoride into tea liquor during infusion. *Environmental Pollution* 104: 197-205.
- FAO/WHO (Food and Agriculture Organization/World Health Organization) Summary and conclusions of the sixty-seventh meeting of the Joint FAO/WHO Expert Committee on Food Additives (JECFA) 2006. [online]
- Grooper SS, Smith JL and Groff JL. (2009) *Advanced Nutrition and Human Metabolism*, Wadsworth, USA: Cengage Learning.
- Hayacibara MF, Queiroz CS, Tabchoury CP, et al. (2004) Fluoride and aluminum in teas and tea-based beverages. *Revista de Saude Publica* 38: 100-105.
- Henderson L, Gregory J and Swan G. (2002) *National Diet and Nutrition Survey: adults aged 19 to 64 years. Volume 1: Types and quantities of foods consumed*, Norwich: The Stationery Office (HMSO).

- 1  
2  
3  
4  
5  
6  
7  
8  
9 Janiszewska J and Balcerzak M. (2013) Analytical problems with the evaluation of  
10 human exposure to fluorides from tea products. *Food Analytical Methods* 6:  
11 1090-1098.
- 12 Jobstl, E., Fairclough, J.P., Davies, A.P., Williamson, M.P. (2005). Creaming in black  
13 tea. *J. Agric. Food Chem*, 53, 7997-8002
- 14 Joshi S, Hlaing T, Whitford GM, et al. (2011) Skeletal fluorosis due to excessive tea  
15 and toothpaste consumption. *Osteoporosis International* 22: 2557-2560.
- 16 Koblar A, Tavčar G and Ponikvar-Svet M. (2012) Fluoride in teas of different types and  
17 forms and the exposure of humans to fluoride with tea and diet. *Food Chemistry*  
18 130: 286-290.
- 19 Koch KR, Pougnet MA, de Villiers S, et al. (1988) Increased urinary excretion of Al  
20 after drinking tea. *Nature* 333: 122.
- 21 Lu Y, Guo WF and Yang XQ. (2004) Fluoride content in tea and its relationship with  
22 tea quality. *Journal of Agricultural and Food Chemistry* 52: 4472-4476.
- 23 Malinowska E, Inkielewicz I, Czarnowski W, et al. (2008) Assessment of fluoride  
24 concentration and daily intake by human from tea and herbal infusions. *Food*  
25 *and Chemical Toxicology* 46: 1055-1061.
- 26 Martínez-Mier EA, Cury J.A., Heilman J.R. , et al. (2011) Development of gold  
27 standard ion-selective electrode-based methods for fluoride analysis *Caries*  
28 *Research* 45: 3-12.
- 29 McArthur A. (2014) Fluoride in tea: Could your daily cuppa really make you ill?  
30 *Nutrition Bulletin* 39: 95-98.
- 31 Medical Research Council. (2002) MRC Working Group report on: Water Fluoridation  
32 and Health. London: MRC.
- 33 Ophaug RH. (1990) Fluoride. In: Brown ML (ed) *Present Knowledge in Nutrition*.  
34 Washington DC: ILSI Press.
- 35 Otten J, Hellwig J and Meyers L. (2006) *Dietary reference intakes: the essential guide*  
36 *to nutrient requirements*, Washington DC: National Academies Press.
- 37 Powell JJ, Greenfield SM, Parkes HG, et al. (1993) Gastro-intestinal availability of  
38 aluminium from tea. *Food and Chemical Toxicology* 31: 449-454.
- 39 Quock RL, Gao JX and Chan JT. (2012) Tea fluoride concentration and the pediatric  
40 patient. *Food Chemistry* 130: 615-617.
- 41 Rao KS and Rao GV. (1994) Effect of aluminium (Al) on brain mitochondrial  
42 monoamine oxidase-A (MAO-A) activity--an in vitro kinetic study. *Molecular*  
43 *and Cellular Biochemistry* 137: 57-60.
- 44 Ruxton CH and Hart VA. (2011) Black tea is not significantly different from water in  
45 the maintenance of normal hydration in human subjects: results from a  
46 randomised controlled trial. *British Journal of Nutrition* 106: 588-595.
- 47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6  
7  
8  
9 Schamschula RG, Sugar E, Un PS, et al. (1985) Physiological indicators of fluoride  
10 exposure and utilization: an epidemiological study. *Community Dentistry &*  
11 *Oral Epidemiology* 13: 104-107.
- 12 Simpson A, Shaw L and Smith AJ. (2001) The bio-availability of fluoride from black  
13 tea. *Journal of Dentistry* 29: 15-21.
- 14 Soto-Rojas AE, Cury J, Heilman J, et al. (2005) Standard Fluoride Analytical Methods  
15 Development: Fluoride analysis of bone by diffusion techniques. *Caries*  
16 *Research* 40: 303.
- 17 Taves DR. (1968) Separation of fluoride by rapid diffusion using hexamethyldisiloxane.  
18 *Talanta* 15: 969-974.
- 19 Tušl J. (1970) Direct determination of fluoride in human urine using fluoride electrode.  
20 *Clinical Chemistry Acta* 27: 216-218.
- 21 Wong MH, Fung KF and Carr HP. (2003) Aluminium and fluoride contents of tea, with  
22 emphasis on brick tea and their health implications. *Toxicology Letters* 137:  
23 111-120.
- 24 Wong MH, Zhang ZQ, Wong JWC, et al. (1998) Trace metal contents (Al, Cu and Zn)  
25 of tea: tea and soil from two tea plantations, and tea products from different  
26 provinces of China. *Environmental geochemistry and health* 20: 87-94.
- 27 World Health Organisation. (1984) Fluorine and Fluorides. *Environmental Health*  
28 *Criteria* 36. Geneva: International Programme on Chemical Safety
- 29 World Health Organization. (1994) Fluorides and oral health. Report of a WHO Expert  
30 Committee on Oral Health Status and Fluoride Use. Geneva, Switzerland.
- 31 World Health Organization. (2010) Inadequate or excess fluoride: a major public health  
32 concern. Geneva, Switzerland.
- 33 Yi J and Cao J. (2008) Tea and fluorosis. *Journal of Fluorine Chemistry* 129: 76-81.
- 34 Yokel RA and Florence RL. (2008) Aluminum bioavailability from tea infusion. *Food*  
35 *and Chemical Toxicology* 46: 3659-3663.
- 36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 1. The effect of Al concentration (mg/l) on the difference between measured F concentration (mg/l) using TISAB III and IV.

Equation for best-fit line:

$$\text{Between-method difference in F concentration (mg/l)} = 0.123 \times \text{Al concentration (mg/l)}$$

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 2. Comparison of mean measured F concentration (mg/l) using TISAB III and IV according to the Al concentration ( $\leq 5$ mg/l and  $> 5$ mg/l) of the samples.

Table 1. Mean (SD) F concentration (mg/l) of 45 tea samples measured at all 3 infusion times plus 2 instant tea samples measured at one infusion time (n=137) using TISAB IV and III and mean (SD) difference in measured F concentration between buffers.

Mean(SD) F concentration (mg/l) measured by using		Difference in measured F concentration (mg/l)*	P value
TISAB IV	TISAB III		
4.37 (2.16)	3.54 (1.65)	0.83 (1.03)	<0.001

\* (TISAB IV- TISAB III)

Table 2. Information on name/brand; form <sup>a</sup> of packaging<sup>a</sup> (B= Tea bag form; L= Loose tea form; I= Instant tea form), caffeine content<sup>b</sup> (C=caffeinated; D= decaffeinated), country of origin and F concentrations (mg/l) at infusion times of 1m, 10m and 1h for 47 black tea samples.

Product name/brand	Tea Form <sup>a</sup> / caffeine content <sup>b</sup>	Country of Origin	F concentration (mg/l) measured by TISAB IV		
			1min	10min	1h
Asda Chosen by You	B/C	Data Unavailable	3.44	6.00	6.32
Lancashire Tea Standard Blend	B/C	Data Unavailable	3.88	5.27	7.30
Morrisons Red Label	B/C	Data Unavailable	3.67	4.41	6.05
PG Tips	B/C	Kenya	6.43	8.41	8.96
Red Label by Sainsbury's	B/C	India, Kenya, Malawi, Sri Lanka	3.19	4.08	5.34
Tesco Original Medium Strength Tea	B/C	Data Unavailable	4.20	6.41	6.86
Tetley Original	B/C	China, India, Sri Lanka	5.40	8.42	8.41
Twinings Original English Breakfast	B/C	Kenya, India, Sri Lanka	1.65	3.23	5.14
Typhoo	B/C	China, Kenya, India	4.64	6.01	6.29
Yorkshire Tea	B/C	Kenya, India	2.16	2.58	6.13
Asda Smart Price Tea Bags	B/C	Data Unavailable	3.23	5.90	8.40
Morrisons Savers	B/C	Data Unavailable	3.33	5.05	5.33
Sainsbury's Basics	B/C	India, Kenya, Malawi, Sri Lanka	4.32	7.07	7.08
Tesco Everyday Value	B/C	Data Unavailable	4.75	7.79	8.95
PG Tips the Fresh One	B/C	Kenya	4.37	6.84	7.06
Twinings the Everyday Tea	B/C	China, India, Kenya, Indonesia	2.98	4.02	5.07
Sainsbury's Taste the Difference Earl Grey	B/C	India, Kenya, Malawi, Sri Lanka	3.65	3.86	3.01
Twinings the Earl Grey	B/C	China	3.01	4.51	4.61
Tetley Earl Grey	B/C	China, India, Sri Lanka	3.51	4.65	5.09
Asda Chosen by You Assam	B/C	India	1.59	2.62	2.74
Asda Chosen by You Ceylon	B/C	Sri Lanka	1.32	1.80	2.07
Tesco Finest Assam	B/C	India	1.28	1.71	1.89
Twinings Assam	B/C	India	1.06	1.38	2.05
Twinings Ceylon	B/C	Sri Lanka	1.20	1.55	1.69
Morrisons Extra Strong	B/C	Data Unavailable	2.96	4.34	4.48
PG Tips the Strong One	B/C	Kenya	5.31	7.40	7.64
Extra Strong Red Label by Sainsbury's	B/C	India, Kenya, Malawi, Sri Lanka	1.96	3.32	3.62
Tetley Extra Strong	B/C	China, India, Sri Lanka	2.12	4.00	4.15



1						
2						
3						
4						
5						
6						
7						
8	Yorkshire Gold	B/C	India, Rwanda	1.44	1.95	2.86
9	Asda Chosen by You					
10	Decaffeinated	B/D	Data Unavailable	2.94	3.00	4.16
11	Morrisons Decaffeinated	B/D	Data Unavailable	3.16	5.52	6.05
12	PG Tips Decaf	B/D	Kenya	6.77	9.28	9.44
13	Decaffeinated Red Label by					
14	Sainsbury's	B/D	India, Kenya, Malawi, Sri Lanka	2.72	5.05	5.25
15	Tesco Decaff Medium Strength					
16	Tea	B/D	Data Unavailable	3.04	4.75	5.80
17	Tetley Decaf	B/D	China, India, Sri Lanka	4.99	7.69	8.66
18	Typhoo Decaf	B/D	China, Kenya, India	3.25	4.55	5.33
19	Yorkshire Tea Decaf	B/D	India, Kenya	3.06	7.72	9.01
20	Asda Chosen by You Loose Leaf					
21	Tea	L/C	Data Unavailable	2.42	2.96	2.82
22	Morrisons Red Label Loose Tea	L/C	Data Unavailable	2.65	3.11	3.25
23	PG Tips Loose Tea	L/C	Kenya	3.92	4.79	4.56
24	Red Label Loose Tea by					
25	Sainsbury's	L/C	India, Kenya, Malawi, Sri Lanka	2.16	2.48	2.66
26	Tesco Original Loose Leaf	L/C	Data Unavailable	3.03	3.66	4.28
27	Twinings Original English					
28	Breakfast Loose	L/C	India, Kenya, Sri Lanka	1.46	1.86	1.62
29	Typhoo Leaf Tea	L/C	China, Kenya, India	1.88	2.29	2.34
30	Yorkshire Tea Leaf Tea	L/C	India, Kenya	2.04	2.51	3.03
31	PG Tips Pure Tea Granules	I/C	Kenya	7.78		
32	Tetley Pure Tea Granules	I/C	China, India, Sri Lanka	9.91		
33						
34						
35						
36						
37						
38						
39						
40						
41						
42						
43						
44						
45						
46						
47						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						

Table 3 Median (range), and mean (SD) F concentration (mg/l) of teas (n=45; excluding instant tea) at 3 infusion times; 1min, 10min and 1h using TISAB IV.

Infusion time	F concentration (mg/l)				
	Median	Minimum	Maximum	Mean	SD
1 m (n=45)	3.04	1.06	6.77	3.16 <sup>a</sup>	1.36
10 m (n=45)	4.41	1.38	9.28	4.57 <sup>a</sup>	2.11
1 h (n=45)	5.14	1.62	9.44	5.17 <sup>a</sup>	2.27
All (n=135)	3.99	1.06	9.44	4.29	2.12

<sup>a</sup> statistically significant difference  $p < 0.001$  (Kruskal-Wallis test, pairwise multiple comparisons analysis)

Table 4. Median (range), and mean (SD) F concentration (mg/l) of 45 tea samples (excluding the 2 instant teas) based on caffeine content and tea form across all 3 infusion times (total number of samples =135).

Sample category	F concentration (mg/l)					
	Median	Minimum	Maximum	Mean	SD	Mann Whitney U
Caffeine content:						
Caffeinated (n=111)	3.93	1.06	8.96	4.04	1.80	P<0.05
Decaffeinated (n=24)	4.72	2.72	9.44	5.47	1.74	
Tea Form:						
Loose tea (n=24)	2.63	1.65	4.42	2.82	0.87	P<0.001
Teabags (n=111)	4.42	1.48	8.50	4.62	1.86	