

Newcastle University ePrints

Farràs P, Benniston AC. [Bodipy-ruthenium\(II\) tris-Bipyridyl Dyads for Homogeneous Photochemical Oxidations.](#)
Tetrahedron Letters 2014, 55(51), 7011-7014.

Copyright:

NOTICE: this is the authors' version of a work that was accepted for publication in *Tetrahedron Letters*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Tetrahedron Letters*, vol. 55, issue 51, 17 December 2014.
DOI: 10.1016/j.tetlet.2014.10.119

Link to published article:

<http://dx.doi.org/10.1016/j.tetlet.2014.10.119>

Date deposited: 13th November 2014



This work is licensed under a [Creative Commons Attribution-NonCommercial 3.0 Unported License](#)

ePrints – Newcastle University ePrints

<http://eprint.ncl.ac.uk>

Bodipy-ruthenium(II) tris-bipyridyl dyads for homogeneous photochemical oxidations

Pau Farràs,* Andrew C. Benniston*

Molecular Photonics Laboratory, School of Chemistry, Newcastle University, Newcastle upon Tyne, NE1 7RU, UK.

Abstract

Two Bodipy-ruthenium(II) tris-bipyridyl dyads were synthesized for use as sensitizers in photochemical oxidation reactions of organic substrates. The synthetic strategy involved the use of a simple “click” CuAAC reaction to link a Bodipy subunit with an organometallic ruthenium(II) tris-bipyridyl complex. The linking triazole bridge was used to minimize electronic coupling between the two subunits. The dyads showed improved performance on organic substrate photo-oxidation reactions compared to the control compound without the Bodipy moiety.

The production of fuels from sunlight remains as one of the main challenges for the scientific community.¹ In recent years, various steps towards efficient systems have been taken, although no example having the desired properties of stability and economic viability has been discovered.² The main components of a full device for light-driven water splitting can be divided as catalysts for oxidation and reduction reactions, light-harvesting compounds, semiconducting electrodes for electron and hole transport and a membrane which both conducts protons and separates the generated gases. Many examples of molecular catalysts have appeared in the literature for water oxidation and proton reduction, some of them exhibiting incredible performances.³ However, there are not many examples of sensitizers for water oxidation, limited mainly by the high oxidation potential needed to drive the reaction and their stability in aqueous media. Common molecular sensitizers for water splitting found in the literature are ruthenium(II) tris(bipyridine) and porphyrin derivatives.⁴ In both cases, the molecules are directly attached to a semiconductor and fast recombination reactions from the injected electrons are observed.⁵


On the other hand, the use of 4,4-difluoro-4-bora-3a,4a-diaza-indacene (Bodipy) as a chromophore has become very popular owing to its exceptional optical properties, photostability, and ease of preparation and purification. Very complicated structures have been prepared based on dendritic scaffolds,⁶ cassette light-harvesters⁷ or multi-dyads showing the versatility of such dyes.⁸ They can be easily functionalised to tune redox and optical properties to meet the demands for the water oxidation reaction. Recently, Bodipy dyes have been published as sensitizers for light-driven hydrogen production although no examples in photochemical oxidation reactions have appeared so far.⁹

In view of such considerations, we envisioned the possibility of designing Bodipy-ruthenium dyads with directional control of electron and hole transport within the sensitizer to minimise recombination reactions and improve the performance of the system. Both subunits are bridged by a triazole linker to avoid coupling between the Bodipy and the ruthenium complex. Moreover, dyad **BDR2** has an ethylene-bipyridyl ligand that is easier to reduce than bipyridine and should direct the electron towards this end of the molecule. The two targeted molecules are shown in Scheme 1 along with the synthetic procedure.

The preparation of dyads **BDR1** and **BDR2** required the synthesis of the new bipyridyl ligand **BDB** (Scheme 1). The starting material azidophenyl-Bodipy (**1**) represents a good candidate to introduce Bodipy fluorophores in organometallic complexes given the reactivity of the azido group.¹⁰ The key step for the preparation of the ligand relies on the flexibility and selectivity of the “click” CuAAC coupling reaction. This

synthetic modular approach opens up the preparation of a large variety of dyads with different types of organic chromophores. Coupling of **1** with **2** under “click” conditions afforded **BDB** in good yield (63%) after careful chromatography (Al₂O₃, eluent: CH₂Cl₂/petroleum ether, 1:1 then 1:0) as a red crystalline material. The ¹H NMR spectrum for **BDB** displayed, along with signals for the Bodipy and bpy groups, a typical singlet at δ 8.63 corresponding to the triazole group.

The ligand **BDB** was complexed using the classical procedure of reacting it with stoichiometric amounts of a metal precursor. The synthetic route used either [Ru^{II}Cl₂(2,2'-bipyridine)₂] **3** or the phenylacetylene derivative **4** as metal salts under ethanolic aqueous solution reflux conditions. The purple precursor **4** was prepared by the reaction of [Ru^{II}Cl₂(DMSO)₂(2,2'-bipyridine)]



Scheme 1. Reagents and conditions: i) CuSO₄, sodium ascorbate, CH₃CN/H₂O, 40 °C, 18 h; ii) EtOH/H₂O, reflux, 18 h, KPF₆(aq). Note: for simplicity the R group for **BDR2** is placed in an arbitrary position.

with **2** in hot DMF. It should be noted that the complex is a mixture of diastereoisomers arising because of the relative positioning of the phenylacetylene group with respect to the chloride ligands and the possibility of Δ and Λ isomers. No attempt was made to separate the diastereoisomers, and so the final dyad **BDR2** is also a mixture of stereoisomers.

All complexes were isolated as the hexafluorophosphate salts by column chromatography and recrystallization from adequate solvents. Their molecular structures were unambiguously assigned by using ^1H , ^{13}C , ^{19}F and ^{11}B NMR spectroscopy, electrospray mass spectrometry and elemental analysis. Both dyads analysed by ESI-MS gave an intense molecular peak with the expected isotopic profile corresponding to the loss of two PF_6^- counter anions assigned to a doubly charged species.

The UV/visible absorption bands of the free ligand and dyad complexes were collected in CH_3CN (Table 1, Figure 1). The absorption and fluorescence profile for the free ligand is typical for a Bodipy derivative.¹¹ The electronic absorption spectrum for **BDB** is dominated by a sharp band centred at 523 nm ($\epsilon_{\text{MAX}} = 75000 \text{ M}^{-1} \text{ cm}^{-1}$) which is assigned to the lowest-energy spin-allowed $\pi\text{-}\pi^*$ transitions involving the Bodipy moiety. The shoulder at the higher energy side of this intense absorption is typical of the vibronic sequence for a dipyrromethene framework. The other weaker transition at *ca.* 380–400 nm is due to the $\text{S}_0 \rightarrow \text{S}_2$ transition. The basic fluorescence spectrum ($\lambda_{\text{FLU}} = 541 \text{ nm}$) at room temperature is on a par with a simple Bodipy derivative, along with the luminescence quantum yield $\Phi_{\text{LUM}} = 0.66$. There is no evidence to suggest that the appended bpy ligand affects the Bodipy group.

In the case for dyads **BDR1** and **BDR2**, the presence of the metal centre is seen by the appearance of bands between 400 and 450 nm. These new bands correspond to the metal-to-ligand charge-transfer (MLCT) transitions between the ruthenium(II) centre and the bpy ligands. For **BDR2**, where the metal centre is surrounded by three different bpy ligands, more electronic transitions arise as evidenced by the presence of a broader band in the region. For both **BDR1** and **BDR2** there is a good match between the observed absorption spectrum and a summation of the individual components (see Supporting Information). Any interaction between the Bodipy and ruthenium complex in the ground state is weak. In comparison to the free ligand, the complexes **BDR1** and **BDR2** give much weaker fluorescence associated with the Bodipy moiety (Figure 1a). The additional band observed at 630 nm (Figure 1b) is typical for Ru(bpy)-type complexes. The total luminescence quantum yield (Φ_{LUM}) (a + b) for both complexes is around 0.03. Preliminary femtosecond

pump-probe experiments indicate that singlet-to-triplet energy transfer in DMF from the excited Bodipy to the Ru(II) component is fast ($\sim 160 \text{ ps}$). Nanosecond flash photolysis results support population of the Bodipy triplet state because of slower ($> 1 \text{ ns}$) reverse energy transfer from the metal-centred $^3\text{MLCT}$ state. A more detailed description of the excited state behaviour will be reported in a full paper.




Figure 1. Absorption (black) and emission (red) spectra for **BDB** (solid), **BDR1** (dash) and **BDR2** (dot) in acetonitrile. Note: a represents emission from the Bodipy moiety and b represents emission from the ruthenium complex.

Electrochemical experiments were carried out by means of cyclic voltammetry in dry acetonitrile containing 0.1 M tetrabutylammonium hexafluorophosphate as a supporting electrolyte. The recorded redox potentials versus SSCE are presented in Table 1. The electrochemical behaviour for the free ligand **BDB** is dominated by an irreversible wave in the oxidative scan, and a quasi-reversible wave upon reductive scanning. The oxidation peak at +1.09 V is assigned to the formation of a Bodipy⁺ (BD^+) unit; the irreversibility is also known for other polypyridyl-containing Bodipy molecules.^{12,13} The quasi-reversible wave seen at -1.15 V is associated with the addition of one electron to the Bodipy site.

The dyads **BDR1** and **BDR2** display an irreversible wave and a quasi-reversible wave upon oxidative scanning. In the reduction scan two well-defined quasi-reversible waves are seen; an extra wave at more negative values is seen as a shoulder, and also observed in similar Ru-BD dyads reported in the literature.¹³

Table 1. Spectroscopic and photophysical data recorded in CH_3CN at 298 K. Oxidation and reduction potentials of the free ligands and dyad molecules vs. sodium saturated calomel electrode (SSCE) in acetonitrile with 0.1 M tetrabutylammonium hexafluorophosphate as a supporting electrolyte at a scan rate of 0.1 V s^{-1} .

Compound	Absorption λ_{max} , nm (ϵ , $\text{M}^{-1} \text{ cm}^{-1}$)	Fluorescence			
		λ_{FLU} , nm	Φ_{LUM}	$E_{1/2\text{ox}}$	$E_{1/2\text{red}}$
BDB	377 (10800), 402 (8400), 493 (27200), 523 (75000)	541	0.66	1.09 (irrev)	-1.15
BDR1	359 (19000), 399 (16800), 430 (19000), 461 (25400), 493 (29400), 524 (66000)	541	0.03	1.05, 1.28	-1.16, -1.31, -1.54, -1.76
BDR2	359 (24600), 398 (20500), 428 (24500), 462 (31000), 492 (31000), 524 (62000)	541	0.03	1.06, 1.30	-1.20, -1.37, -1.46

Notwithstanding, the electrochemical behaviour for **BDR1** is readily understood as a near perfect overlap of the cyclic voltammograms observed for the two halves of the molecule; namely, ligand **BDB** and $[\text{Ru}(\text{bpy})_3]^{2+}$. This behaviour is observed for both dyad molecules. As an example, the **BDR1** oxidative segment is dominated by an irreversible wave at +1.05

V (Bodipy-based) followed by a reversible one-electron wave at +1.28 V (Ru-based). In the reduction side, once again is seen a wave at -1.16 V (reduction of Bodipy), and three quasi-reversible waves at -1.31, -1.54 and -1.76 V ($\Delta E_p = 80 \text{ mV}$) which are typical for Ru(bpy)-type complexes.¹⁴ From these results it is inferred that the Ru and Bodipy subunits are isolated and only

minor, if any, electronic communication takes place through the bridge in the ground state.

The ability of dyads **BDR1** and **BDR2** to act as sensitizers in the photochemical oxidation of water and organic substrates was tested with two molecular ruthenium oxidation catalysts reported in the literature. Catalyst $[\text{Ru}^{\text{II}}(\text{bda})(4\text{-bromopyridine})_2]$ where ($\text{H}_2\text{bda} = 2,2\text{-bipyridine-6,6-dicarboxylic acid}$), **RuCAT**, is known to be a good water oxidation catalyst under visible light illumination using $[\text{Ru}(4,4\text{-(PO}_3\text{H}_2)_2\text{-bpy})(\text{bpy})_2]^{2+}$ as the sensitizer.¹⁵ However, the latter complex has an oxidation potential of +1.20 V vs. SSCE, which is higher than the oxidation potential of the $\text{BD}^{\text{I/O}}$ couple in the dyads (+1.05 V). Electrochemical analysis by CV of a mixture of **BDR1** and **RuCAT** is shown in Figure 2. The black trace shows the redox couples corresponding to dyad **BDR1** ($\text{BD}^{\text{I/O}}$ and $\text{Ru}_p^{\text{III/II}}$) in a water:acetonitrile mixture.




Figure 2. Redox properties of dyad **BDR1** (black) and **RuCAT** (red) in a water:acetonitrile mixture using a glassy carbon disk electrode, a Pt wire as the auxiliary, and a SSCE as the reference electrode, at a scan rate of 100 mV s^{-1} .

The red trace in Figure 2 shows the formation of the $\text{Ru}_c^{\text{II}}\text{-OH}_2$ species from the water oxidation catalyst **RuCAT** and its higher oxidation states. Upon addition of water, **RuCAT** coordinates an aquo ligand which modulates its redox properties. It is known that for water oxidation to take place the generation of at least a $\text{Ru}_c^{\text{V}}\text{=O}$ species is needed (electrocatalytic wave starting at +1.20 V). Such a species is not achievable for **BDR1** because of its lower oxidation potential. However, the $\text{Ru}_c^{\text{IV}}\text{=O}$ oxidation state can be obtained and, therefore, oxidation of organic substrates should be achieved with **RuCAT** and the Bodipy-Ru dyads. A second catalyst $\{[\text{Ru}^{\text{II}}(\text{tpy})_2(\mu\text{-pyr-dc})(\mu\text{-OOCMe})]\}$ [**RuCAT2**] was also used which is known to be a very good oxidation catalyst for organic substrates.¹⁶

The ability of dyads **BDR1** and **BDR2** and control compound $[\text{Ru}(\text{bpy})_3]^{2+}$ to act as sensitizers in photochemical oxidation reactions were examined for the oxidation of a variety of organic substrates and the results obtained are presented in Table 2. Typically, we used a ratio of 1:5:500:1000 of catalyst:sensitizer:substrate:electron acceptor and a white LED ring lamp (0.05 W cm^{-2}) for 2.5 hours in a thermostated jacketed cell. As is observed in entries 1-4 of Table 2 for $4\text{-OCH}_3\text{-C}_6\text{H}_4\text{CH}_2\text{OH}$, the reaction only proceeds when all components are present in the reaction. In general, the catalytic activity observed when the control compound is used is lower than when using the dyads under the same reaction conditions. Actually, the

best performance in our case was obtained with dyad **BDR1**, that yields 65 TN with a conversion of 13%. This indicates that the presence of the Bodipy subunit in the sensitizer favors the photocatalytic reaction. The same trend is observed in entries 5-7 where **RuCAT2** was used as the oxidation catalyst. In this case, the performance of **RuCAT2** is improved compared to **RuCAT**, as expected for this type of ruthenium catalysts. Other alcohols tested with **BDR1** and **RuCAT2** show a very different reactivity, whereas 2-propanol is easily oxidized to acetone, $3\text{-NO}_2\text{-C}_6\text{H}_4\text{CH}_2\text{OH}$ only yields 25 TN with a conversion of 5%.

Sulfides were also tested as substrates as indicated in entries 11-13, under the same conditions as the previous substrates, giving a good TN of 420 that represents a conversion of 84%. Again, the use of the control compound gave a lower conversion yield of 61% with a 305 TN. Even though sulfides are easier to oxidize than alcohols, a general trend is observed in all cases where Bodipy-Ru dyads perform better than the control compound. However, only a minor difference is observed between both dyads, indicating that the presence of the directional ethylene-bipyridyl ligand does not affect the electron transfer processes within the sensitizer.

Table 2. Photocatalytic oxidation of a variety of substrates using dyads **BDR1** and **BDR2** and control compound $[\text{Ru}(\text{bpy})_3]^{2+}$ in aqueous solution.^a

Entry	Sensitizer	Cat	Substrate	TN (Conv. %)
1	-	RuCAT	$4\text{-OCH}_3\text{-C}_6\text{H}_4\text{CH}_2\text{OH}$	-
2	$[\text{Ru}(\text{bpy})_3]^{2+}$			50 (10)
3	BDR1			65 (13)
4	BDR2			65 (13)
5	$[\text{Ru}(\text{bpy})_3]^{2+}$	RuCAT2	$4\text{-OCH}_3\text{-C}_6\text{H}_4\text{CH}_2\text{OH}$	70 (14)
6	BDR1			100 (20)
7	BDR2			95 (19)
8	BDR1	RuCAT2	$4\text{-Br-C}_6\text{H}_4\text{CH}_2\text{OH}$	100 (20)
9			2-propanol	175 (35)
10			$3\text{-NO}_2\text{-C}_6\text{H}_4\text{CH}_2\text{OH}$	25 (5)
11	$[\text{Ru}(\text{bpy})_3]^{2+}$	RuCAT2	$4\text{-Br-C}_6\text{H}_4\text{SMe}$	305 (61)
12	BDR1			420 (84)
13	BDR2			415 (83)

^a Reaction conditions: photosensitizer (0.1 mM)/catalyst (0.02 mM)/substrate (10 mM)/ $[\text{Co}^{\text{III}}(\text{NH}_3)_5\text{Cl}]^{2+}$ (20 mM) irradiated with a white LED ring lamp (0.05 W cm^{-2}) at 298 K in a borate buffer aqueous solution (pH 7, 20 mM).

In conclusion, we have shown that Bodipy subunits can be readily incorporated within $\text{Ru}(\text{bpy})$ -like sensitizers for photochemical oxidation reactions. In addition, the azidophenyl-Bodipy proved to be a promising starting material for incorporating Bodipy subunits within donor-acceptor dyads by formation of the triazole bridge. We expect that the present

systems can be modified with anchoring groups via the Bodipy or ruthenium complex site for their attachment to semiconducting metal oxides. One intention is to remove the necessity of using a sacrificial agent by coupling two half-redox reactions in a photochemical cell.

Acknowledgments

P.F. gratefully acknowledges the Royal Society for the Newton Fellowship and for the financial support. We also thank the Engineering and Physical Sciences Research Council (EPSRC) sponsored Mass Spectrometry at Swansea for the collection of mass spectra.

Supplementary Material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/...>

References

- (a) Directing Matter and Energy: Five Challenges for Science and the Imagination; U.S. Dept. of Energy: Washington, D. C., December, 2007; (b) Basic Research Needs: Catalysis For Energy; U.S. Dept. of Energy: Washington, D. C., August, 2007.
- (a) Kanan, M. W.; Nocera, D. G. *Science* **2008**, *321*, 1072-1075; (b) Youngblood, W.; Lee, S.; Kobayashi, Y.; Hernandez-Pagan, E.; Hoertz, P.; Moore, T.; Moore, A.; Gust, D.; Mallouk, T. *J. Am. Chem. Soc.* **2009**, *131*, 926-927; (c) Brimblecombe, R.; Koo, A.; Dismukes, G.; Swiegers, G.; Spiccia, L. *J. Am. Chem. Soc.* **2010**, *132*, 2892-289; (d) Li, L.; Duan, L.; Xu, Y.; Gorlov, M.; Hagfeldt, A.; Sun, L. *Chem. Commun.* **2010**, *46*, 7307-7309; (e) Gao, Y.; Ding, X.; Liu, J.; Wang, L.; Lu, Z.; Li, L.; Sun, L. *J. Am. Chem. Soc.* **2013**, *135*, 4219-4222.
- (a) Yin, Q.; Tan, J. M.; Besson, C.; Geletii, Y. V.; Musaev, D. G.; Kuznetsov, A. E.; Luo, Z.; Hardcastle, K. I.; Hill, C. L. *Science* **2010**, *328*, 342-345; (b) Toma, F. M.; Sartorel, A.; Iurlo, M.; Carraro, M.; Parisse, P.; Maccato, C.; Rapino, S.; Gonzalez, B. R.; Amenitsch, H.; Da Ros, T.; Casalis, L.; Goldoni, A.; Marccaccio, M.; Scorrano, G.; Scoles, G.; Paolucci, F.; Prato, M.; Bonchio, M. *Nat. Chem.* **2010**, *2*, 826-831; (c) Fillol, J. L.; Codolà, Z.; Garcia-Bosch, I.; Gómez, L.; Pla, J. J.; Costas, M. *Nat. Chem.* **2011**, *3*, 807-813; (d) Helm, M. L.; Stewart, M. P.; Bullock, R. M.; DuBois, M. R.; DuBois, D. L. *Science* **2011**, *333*, 863-866; (e) Duan, L.; Bozoglian, F.; Mandal, S.; Stewart, B.; Privalov, T.; Llobet, A.; Sun, L. *Nat. Chem.* **2012**, *4*, 418-423.
- (a) Puntoriero, F.; La Ganga, G.; Sartorel, A.; Carraro, M.; Scorrano, G.; Bonchio, M.; Campagna, S. *Chem. Commun.* **2010**, *46*, 4725-4727; (b) Moore, G. F.; Konezny, S. J.; Song, H. E.; Milot, R. L.; Blakemore, J. D.; Lee, M. L.; Batista, V. S.; Schmuttenmaer, C. A.; Crabtree, R. H.; Brudvig, G. W. *J. Phys. Chem. C* **2012**, *116*, 4892-4902.
- (a) Hanson, K.; Torelli, D. A.; Vannucci, A. K.; Brennaman, M. K.; Luo, H. L.; Alibabaei, L.; Song, W. J.; Ashford, D. L.; Norris, M. R.; Glasson, C. R. K.; Concepcion, J. J.; Meyer, T. J. *Angew. Chem. Int. Ed.* **2012**, *51*, 12782-12785; (b) Nayak, A.; Knauf, R. R.; Hanson, K.; Alibabaei, L.; Concepcion, J. J.; Ashford, D. L.; Dempsey, J. L.; Meyer, T. J. *Chem. Sci.* **2014**, *5*, 3115-3119.
- Kostereli, Z.; Ozdemir, T.; Buyukcikir, O.; Akkaya, E. U. *Org. Lett.* **2012**, *14*, 3636-3639.
- Ziessel, R.; Ulrich, G.; Haefele, A.; Harriman, A. *J. Am. Chem. Soc.* **2013**, *135*, 11330-11344.
- Bandi, V.; Das, S. K.; Awuah, S. G.; You, Y.; D'Souza, F. *J. Am. Chem. Soc.* **2014**, *136*, 7571-7574.
- (a) Sabatini, R. P.; McCormick, T. M.; Lazarides, T.; Wilson, K. C.; Eisenberg, R.; McCamant, D. W. *J. Phys. Chem. Lett.* **2011**, *2*, 223-227; (b) Bartelmess, J.; Francis, A. J.; El Roz, K. A.; Castellano, F. N.; Weare, W. W.; Sommer, R. D. *Inorg. Chem.* **2014**, *53*, 4527-4534.
- (a) Bai, D.; Benniston, A. C.; Hagon, J.; Lemmetyinen, H.; Tkachenko, N. V.; Clegg, W.; Harrington, R. W. *Phys. Chem. Chem. Phys.* **2012**, *14*, 4447; (b) Bai, D.; Benniston, A. C.; Hagon, J.; Lemmetyinen, H.; Tkachenko, N. V.; Harrington, R. W. *Phys. Chem. Chem. Phys.* **2013**, *15*, 9854; (c) Bai, D.; Benniston, A. C.; Whittle, V. L.; Lemmetyinen, H.; Tkachenko, N. V. *ChemPhysChem* **2014**, *15*, 3089.
- Benniston, A. C.; Copley, G. *Phys. Chem. Chem. Phys.* **2009**, *11*, 4124.
- Galletta, M.; Campagna, S.; Quesada, M.; Ulrich, G.; Ziessel, R. *Chem. Commun.*, **2005**, 4222.
- Galletta, M.; Puntoriero, F.; Campagna, S.; Chiorboli, C.; Quesada, M.; Goeb, S.; Ziessel, R. *J. Phys. Chem. A* **2006**, *110*, 4348.
- Juris, A.; Balzani, V.; Barigelletti, F.; Campagna, S.; Belser, P., von Zelewsky, A. *Coord. Chem. Rev.* **1988**, *84*, 85-277.
- Duan, L.; Wang, L.; Inge, A. K.; Fischer, A.; Zou, X.; Sun, L. *Inorg. Chem.* **2013**, *52*, 7844-7852.
- Di Giovanni, C.; Poater, A.; Benet-Buchholz, J.; Cavallo, L.; Solà, M.; Llobet, A. *Chem. Eur. J.* **2014**, *20*, 3898-3902.