
Tan KH, Sheng KY, Chang CY, Yin WF, Chan KG. [Draft genome sequence of a quorum-sensing bacterium, *Dickeya* sp. strain 2B12, isolated from a freshwater lake](#). *Genome Announcements* 2015, 3(1), e01542-14.

Copyright:

© 2015 Tan et al.

This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](#).

DOI link to article:

<http://dx.doi.org/10.1128/genomeA.01542-14>

Date deposited:

04/01/2016



This work is licensed under a [Creative Commons Attribution 3.0 Unported License](#)

Draft Genome Sequence of a Quorum-Sensing Bacterium, *Dickeya* sp. Strain 2B12, Isolated from a Freshwater Lake

Kian-Hin Tan,^a Kit-Yeng Sheng,^a Chien-Yi Chang,^{b,c} Wai-Fong Yin,^a  Kok-Gan Chan^a

Division of Genetics and Molecular Biology, Institute of Biological Sciences, Faculty of Science, University of Malaya, Kuala Lumpur, Malaysia^a; Interdisciplinary Computing and Complex BioSystems (ICOS) research group, School of Computing Science, Newcastle University, Newcastle upon Tyne, United Kingdom^b; The Centre for Bacterial Cell Biology, Medical School, Newcastle University, Newcastle upon Tyne, United Kingdom^c

***Dickeya* sp. strain 2B12 was isolated from a freshwater lake in Malaysia. Here, we report the draft genome sequence of *Dickeya* sp. 2B12 sequenced by the Illumina MiSeq platform. With the genome sequence available, this genome sequence will be useful for the study of quorum-sensing activity in this isolate.**

Received 19 December 2014 Accepted 22 December 2014 Published 5 February 2015

Citation Tan K-H, Sheng K-Y, Chang C-Y, Yin W-F, Chan K-G. 2015. Draft genome sequence of a quorum-sensing bacterium, *Dickeya* sp. strain 2B12, isolated from a freshwater lake. *Genome Announc* 3(1):e01542-14. doi:10.1128/genomeA.01542-14.

Copyright © 2015 Tan et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Kok-Gan Chan, kokgan@um.edu.my.

All members of the *Enterobacteriaceae* family that are pathogenic to plants, both pectolytic (*Erwinia carotovora* and *Erwinia chrysanthemi*) and nonpectolytic (*Erwinia amylovora*), were assigned into a new genus, *Erwinia*, in 1917 (1). Despite the original notion that *Erwinia chrysanthemi* is a pathogen of chrysanthemum (2), and thus named as such, it was later discovered to infect a wide range of plant hosts (3, 4). *E. chrysanthemi* strains were classified into several pathovars according to pathogenicity on host plants and biochemical and physiological differences (5). This species was later transferred into a new genus called *Dickeya*, which now comprises eight species (*D. chrysanthemi*, *D. dadantii*, *D. dianthicola*, *D. dieffenbachiae*, *D. zaeae*, *D. paradisiaca*, *D. solani*, and *D. aquatica*) (1, 2, 6). Here, we present a draft genome sequence of *Dickeya* sp. strain 2B12, isolated from a fresh lake water sample.

The genome of *Dickeya* sp. 2B12 was sequenced by an Illumina MiSeq sequencer with a 150-bp read chemistry. In brief, genomic DNA of *Dickeya* sp. 2B12 was extracted from an overnight culture using a Master Pure DNA purification kit (Epicentre, Inc., Madison, WI, USA). The quality and quantity of the DNA was assessed via a NanoDrop spectrophotometer (ThermoScientific, Waltham, MA, USA) and a Qubit 2.0 fluorometer (Life Technologies, Carlsbad, CA, USA). Then, 50 ng of DNA was used to construct a sequencing library using a Nextera DNA sample prep kit (Illumina, San Diego, CA, USA) according to the manufacturer's protocol. The library was quantified by a Qubit 2.0 fluorometer and Bioanalyzer high-sensitivity chip (Agilent Technologies, Santa Clara, CA, USA). A sequencing run was set up to generate 2 × 150 base-paired reads.

The raw reads were trimmed at Q30, resulting in 1,745,868 reads with an average length of 144.3 bp. The reads were assembled *de novo* using CLC Genomic Workbench 6 (CLC Bio, Denmark), giving 120 contigs with an N_{50} of 92,830 bp, constituting a genome size of 4,349,822 bp. The assembled genome has a GC content of 54.5%. There are 107 contigs with an average coverage of 30×, and the average coverage across the genome is 46.7×. This

genome was annotated using Rapid Annotation using Subsystem Technology (RAST), version 2.0 (7), which revealed the presence of 3,965 coding sequences and 77 RNA genes, with 59% of the coding sequences (CDS) covered by the subsystem in the RAST server. The *Dickeya* sp. 2B12 harbors putative pectate lyase and cellulase genes in its genome, which suggest the potential of *Dickeya* sp. 2B12 as a plant pathogen, a well-known role performed by other members of *Dickeya* spp. (1, 2). Interestingly, a pair of *luxI*/*luxR* homologues was also present in the genome in contig 11, suggesting the adaptation of a quorum-sensing system to regulate gene expression in this bacterium.

Nucleotide sequence accession number. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [JSYG000000000](https://www.ncbi.nlm.nih.gov/nuccore/JSYG000000000). The version described in this paper is the first version.

ACKNOWLEDGMENTS

This research was financed by University of Malaya High-Impact Research (HIR) grants (UMC/625/1/HIR/MOHE/CHAN/01 [no. A-000001-50001] and UM-MOHE HIR UMC/625/1/HIR/MOHE/CHAN/14/1 [no. H-50001-A000027]) (to Kok-Gan Chan), which we gratefully acknowledge.

REFERENCES

- Toth IK, van der Wolf JM, Saddler G, Lojkowska E, Hélias V, Pirhonen M, Tsror L, Elphinstone JG. 2011. *Dickeya* species: an emerging problem for potato production in Europe. *Plant Pathol* 60:385–399. <http://dx.doi.org/10.1111/j.1365-3059.2011.02427.x>.
- Burkholder WD, McFadden LA, Dimock AW. 1953. A bacterial blight of chrysanthemums. *Phytopathology* 43:522–526.
- Samson R, Legendre JB, Christen R, Fischer-Le Saux M, Achouak W, Gardan L. 2005. Transfer of *Pectobacterium chrysanthemi* (Burkholder *et al.* 1953) Brenner *et al.* 1973 and *Brenneria paradisiaca* to the genus *Dickeya* gen. nov. as *Dickeya chrysanthemi* comb. nov. and *Dickeya paradisiaca* comb. nov. and delineation of four novel species, *Dickeya dadantii* sp. nov., *Dickeya dianthicola* sp. nov., *Dickeya dieffenbachiae* sp. nov. and *Dickeya zaeae* sp. nov. *Int J Syst Evol Microbiol* 55:1415–1427. <http://dx.doi.org/10.1099/ijs.0.02791-0>.
- Ma B, Hibbing ME, Kim HS, Reedy RM, Yedidia I, Breuer J, Breuer J,

- Glasner JD, Perna NT, Kelman A, Charkowski AO. 2007. Host range and molecular phylogenies of the soft rot enterobacterial genera *Pectobacterium* and *Dickeya*. *Phytopathology* 97:1150–1163. <http://dx.doi.org/10.1094/PHYTO-97-9-1150>.
5. Dye DW, Bradbury JF, Goto M, Hayward AC, Lelliott RA, Schroth MN. 1980. International standards for naming pathovars of phytopathogenic bacteria and a list of pathovar names and pathotype strains. *Rev Plant Pathol* 59:59153–59168.
6. Parkinson N, DeVos P, Pirhonen M, Elphinstone J. 2014. *Dickeya aquatica* sp. nov., isolated from waterways. *Int J Syst Evol Microbiol* 64: 2264–2266. <http://dx.doi.org/10.1099/ijs.0.058693-0>.
7. Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res* 42: D206–D214. <http://dx.doi.org/10.1093/nar/gkt1226>.