

Ross A, Yarnall AJ, Rochester L, Lord S. [A novel approach to falls classification in Parkinson's disease: Development of the Fall-Related Activity Classification \(FRAC\)](#). *Physiotherapy* 2016

Copyright:

©2016. This manuscript version is made available under the [CC-BY-NC-ND 4.0 license](#)

DOI link to article:

<http://dx.doi.org/10.1016/j.physio.2016.08.002>

Date deposited:

19/10/2016

Embargo release date:

03 September 2017



This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International licence](#)

1 A novel approach to falls classification in Parkinson's disease: Development of the Fall-
2 Related Activity Classification (FRAC)

3 Annie Ross¹, Alison J Yarnall¹, Lynn Rochester¹, Sue Lord* ¹

4

5 ¹Institute of Neuroscience, Newcastle University Institute for Aging, Newcastle upon Tyne,
6 UK

7

8 **Correspondence to:**

9 Sue Lord PhD

10 Institute of Neuroscience,

11 Newcastle University Institute for Aging,

12 Newcastle University

13 Newcastle upon Tyne

14 NE4 5PL

15 Email: sue.lord@ncl.ac.uk

16

17 **Running title:** A novel classification for falls

18 **Key Words:** falls, classification, reliability

19 **Manuscript word count:** 2400

20

21 * Corresponding author.

22 Institute of Neuroscience, Newcastle University, Clinical Ageing Research Unit, Campus for
23 Ageing and Vitality, Newcastle upon Tyne NE4 5PL, UK. Tel.: +44(0)191 2081291.

24

25 E-mail addresses: annie.ross@nuth.nhs.uk (A. Ross), alison.yarnall@newcastle.ac.uk (A.

26 Yarnall), lynn.rochester@newcastle.ac.uk (L. Rochester), sue.lord@newcastle.ac.uk (S. Lord)

27

28

29

30

31 **ABSTRACT**

32 **Background:** Falls are a major problem for people with Parkinson's disease (PD). Despite
33 years of focused research knowledge of falls aetiology is poor. This may be partly due to
34 classification approaches which conventionally report fall frequency. This nosology is blunt,
35 and does not take into account causality or the circumstances in which the fall occurred. For
36 example, it is likely that people who fall from a postural transition are phenotypically
37 different to those who fall during high level activities. Recent evidence supports the use of
38 a novel falls classification based on fall related activity, however its clinimetric properties
39 have not yet been tested.

40 **Objective:** This study describes further development of the Fall-Related Activity
41 Classification (FRAC) and reports on its inter-rater reliability (IRR).

42 **Method:** Descriptors of the FRAC were refined through an iterative process with a
43 multidisciplinary team. Three categories based on the activity preceding the fall were
44 identified. PD fallers were categorised as: 1) advanced 2) combined or 3) transitional. Fifty-
45 five fall scenarios were rated by 23 raters using a standardised process. Raters comprised 3
46 clinical subgroups: 1) physiotherapists, 2) physicians, 3) non-medical researchers. IRR
47 analysis was performed using weighted kappa coefficients and included sub group analysis
48 based on clinical speciality.

49 **Results:** Excellent agreement was reached for all clinicians, $\kappa = 0.807$ (95%CI 0.732-0.870).
50 Clinical subgroups performed similarly well (range of $\kappa = 0.780 - 0.822$).

51 **Conclusion:** The FRAC can be reliably used to classify falls. This may discriminate between
52 phenotypically different fallers and subsequently strengthen falls predictors in future
53 studies.

54 **INTRODUCTION**

55 Falls are a major problem in Parkinson's disease (PD) with a recent systematic review
56 demonstrating that 60% of people with PD fall at least once each year with 39% reporting
57 recurrent falls [1]. Many studies have investigated risk factors for falls [2-5] yet our
58 understanding of falls predictors remain limited. Falls are multifactorial with a complex
59 pathophysiology [6] and solely reporting frequency of falls may not be sensitive enough to
60 accurately identify risk factors. The most robust prediction of falls is two or more falls in the
61 previous year [4], however there is also a clinical need to identify prodromal fallers in order
62 to prevent the spiral of fear of falling and reduced physical activity with consequent
63 functional decline [7].

64

65 Conventionally in falls research, falls frequency is used as the primary outcome without
66 consideration of the circumstances in which the fall occurred. Understanding this is
67 important because it is likely that people who fall whilst turning or standing up are
68 phenotypically different to those who fall during higher level activities such as while walking
69 [5]. Different risk factors may therefore underpin falls risk; a distinction that is lost when
70 reporting frequency alone. Although falls frequency is sensitive to risk factors such as
71 disease severity and recurrent fallers show different fall characteristics and clinical features
72 to single fallers [8], this knowledge has limited clinical utility. Importantly, established
73 (recurrent) falls are challenging to manage effectively, with recent evidence suggesting that

74 interventions to reduce falls are more effective for people with mild disease severity [9, 10].
75 Ideally, interventions will occur *prior* to the first fall occurring. This requires a more nuanced
76 approach to assessment and classification, and recognition of different faller phenotypes.

77

78 Earlier work has examined non-frequency based classifications. For example, fall related
79 activity has been described in relation to falls [5, 8, 11, 12] with walking identified as the
80 most common fall related activity [11, 12]. Location has also been shown to influence risk
81 factors [13, 14] with indoor falls associated with disability, poor health and inactivity, in
82 contrast to outdoor falls which are associated with an active lifestyle and average or better
83 than average health [13]. Another approach is to describe the mechanism of the fall [15-18]
84 using terms such as 'extrinsic' which classifies the circumstance surrounding the fall and
85 may include environmental descriptors (e.g. obstacle, hazards) or the specific fall related
86 activity. Other descriptors include biomechanical perturbation that preceded the fall [19,
87 20] for example a 'base of support' fall (a slip or trip), or a 'centre of mass' fall (bending or
88 reaching). Very few studies have categorised falls based on fall related activity such as: a)
89 transferring, stooping, bending, or standing still; (b) walking; (c) turning around or reaching;
90 (d) going up or down stairs, steps or curbs; and (e) "high risk" activities like running or
91 standing on a chair [12, 21]. However, apart from one classification which reported a
92 reliability of $\kappa = 0.828$ [17], none of these classifications have been formally scripted, tested
93 or adopted.

94

95 We considered the advantages and disadvantages of previous falls classifications, and also
96 conducted some preliminary research that resulted in our decision to adopt a novel
97 approach. For the preliminary research, we explored the relationship between ambulatory

98 activity and falls in people with PD and compared this association for falls frequency and
99 falls context. In order to do this, we first developed a classification based on fall related
100 activity. At 12 months 36.9% of the cohort had fallen. Total time spent walking was
101 significantly lower for transitional fallers compared with non-fallers and they also had
102 significantly increased disease severity. There were no significant relationships when fallers
103 were categorized by frequency. This demonstrates greater discrimination for fallers versus
104 non-fallers when the falls context classification was used [5]. However, reliability testing of
105 the classification was limited to an informal assessment whereby four raters classified 20 fall
106 scenarios from which a Fleiss' kappa coefficient $\kappa = 0.643$ (95% CI 0.513-0.686) was obtained.
107 The aim of this study was therefore to formally examine the inter-rater reliability of this falls
108 context classification, which we named the Fall-Related Activity Classification (FRAC) [5]. We
109 also examined reliability results for raters clustered by clinical and falls expertise because
110 we were interested in its generalisability.

111

112 **METHODS**

113 *Description of Fall-Related Activity Classification (FRAC)*

114 The original definitions and descriptors of the FRAC [5] were reworked and the original title,
115 "ambulatory" was renamed "combined". Three categories are described based on a
116 continuum of everyday activities (see Table 1 and Figure 1).

117 *Reliability study fall scenarios*

118 The fall scenarios for this reliability study were taken from the first 12 months of falls diaries
119 from the ICICLE-Gait study falls database.

120 This is a collaborative study with ICICLE-PD, an incident cohort study (Incidence of Cognitive
121 Impairment in Cohorts with Longitudinal Evaluation - Parkinson's disease); full description of
122 this cohort is available elsewhere [22]. Briefly, the authors aimed to recruit all cases of
123 incident idiopathic PD from secondary care services in Newcastle-upon-Tyne and Gateshead
124 between June 2009 and December 2011. ICICLE-GAIT recruited a subset of the cohort at the
125 same time point. Primary care (general practitioners) and secondary care (neurologists,
126 geriatricians and PD specialist nurses) services were invited to notify the investigators of
127 potential participants. Participants had their PD diagnosis confirmed by a consultant
128 neurologist specialising in neurodegenerative diseases according to the UK Brain Bank
129 Criteria [23]. Exclusion criteria included a diagnosis of Parkinsonism prior to study onset and
130 non-idiopathic forms of the disease, such as drug-induced and vascular Parkinsonism and
131 the atypical Parkinsonism syndromes including supranuclear palsy, multiple system atrophy
132 or cortico-basal degeneration. This was to ensure that only cases of incident idiopathic PD
133 were included. Participants were also excluded if they had evidence of a significant memory
134 impairment or dementia, as evidenced by a Mini Mental State Examination (MMSE) score
135 <24 or did not have sufficient knowledge of the English language in order to co-operate with
136 testing. The study was approved by the Newcastle and North Tyneside Research Ethics
137 Committee and all participants gave informed consent.

138

139 Throughout the study period participants were asked to record any falls that occurred in the
140 past month on a standardised prompt sheet, including the date and time of each fall as well
141 as location, preceding activity, perceived cause, position in which they landed and mode of
142 recovery in a structured open-ended statement format. All reported falls were followed-up
143 with a telephone call from a Senior Research Physiotherapist (DM) to verify information and

144 rectify any missing data. Fifty five falls scenarios were randomly selected from the first 12
145 months of recorded falls.

146

147 *Procedure*

148 A convenience sample of 25 raters from gerontology and neurology clinical and research
149 backgrounds within the local NHS Trust agreed to participate in the study. Standardisation
150 of the rating process was established to prevent bias. Raters were instructed to read
151 through the definitions and examples, familiarise themselves with the FRAC and then
152 independently categorise 55 scenarios in to one of three categories (see Table 1). No
153 formal training was provided. Raters were blinded to any information other than the
154 necessary details regarding the fall related activity needed to classify. Raters were asked 4
155 questions about their clinical background in order to answer the secondary research
156 question. An “expert in falls” was defined as those who work regularly/have worked
157 regularly in the past in falls clinics, or those who routinely assess and treat older patients at
158 risk of falls or following a fall in their clinical practice [24].

159

160 *Statistical analysis*

161 In order to minimise bias this study incorporated a fully crossed design, meaning that all falls
162 scenarios were rated by all raters. Light’s [25] kappa was used to assess IRR because there
163 were more than 3 raters [26]. Squared weighting of errors was used to give partial credit for
164 judgements that disagree but are close. The “psy” package of R statistics v3.1.0 software
165 was used to calculate the kappa values and 95% confidence intervals were calculated using
166 the “boot” package.

167

168 **RESULTS**

169 *Sample characteristics of raters*

170 The final number of raters was 23. Eleven raters were physiotherapists, seven were
171 physicians and five were non-medical researchers (bioengineer, bio mechanist, engineer,
172 researcher and research technician). The mean number of years of experience working
173 specifically with people with PD was 3.99 (range 0- >15 years), 16 raters had regular
174 experience with fallers and six raters were deemed to be “falls experts”.

175

176 *Distribution of types of falls rated*

177 With respect to the most frequently chosen response across the 23 raters, 16 falls were
178 classified as transitional, 27 were classified as combined and 12 were classified as advanced.
179 Some more complex falls scenarios resulted in poor agreement (see Table 2).

180

181 *Statistical estimates of inter-rater reliability with regards to rater population*

182 Subgroups of clinicians performed similarly with all values indicating substantial agreement
183 [27] and there was no statistically significant difference between the groups (see Table 3).

184

185 *Statistical estimates of inter-rater reliability*

186 Kappa was computed for each coder pair then averaged to provide a single index of IRR [25].
187 Excellent agreement was reached $\kappa = 0.807$ (95%CI 0.732-0.870) (see Table 3).

188

189

190 **DISCUSSION**

191 This study reports excellent inter-rater reliability for the FRAC suggesting it is a robust
192 classification to identify falls. Importantly, reliability was upheld for all raters including non-
193 medical clinicians and raters with fewer years of experience in falls management. Further
194 work is required to validate the classification and more extensively test its clinimetric
195 properties. Although the FRAC was developed for people with PD its application is likely to
196 be broader. A fall is a generic event, and there is no reason why this classification should be
197 limited to rating by a specific discipline. Future research will examine the utility and
198 clinimetric properties of the FRAC for other populations including older adults, and
199 investigate acceptability and utility.

200

201 Comparison with earlier reliability studies is limited, however results are in line with the
202 only other falls classification to report reliability [17]. In this study two reviewers classified
203 falls from a sample of community-dwelling adults into one of four categories and
204 demonstrated $k = 0.828$. The taxonomy was extensive with four major categories each
205 encompassing three levels: extrinsic (including falls, slips, trips); intrinsic (including mobility
206 or balance disorders); falls from a non-bipedal stance (such as falls from bed/chair); and
207 unclassifiable falls. This classification is limited in its clinical applicability and has not been
208 widely adopted. In addition, only two raters classified falls using this taxonomy and both
209 were geriatric nurses. It has been suggested that studies that include ratings from a single
210 professional group suffer the risk that professional assumptions underlie the ratings, which
211 may artificially inflate the reliability of the instrument [28]. In contrast, 23 raters from a
212 variety of professional backgrounds validated the FRAC.

213

214 The FRAC is relatively simple with only three categories to select from and includes concise
215 descriptors and a written guide to assist with classification. Also, no formal training is
216 required to use it reliably, which demonstrates generalisability. The key limitation we noted
217 was that some of the more complex falls scenarios resulted in poorer agreement because it
218 was challenging to interpret the level of complexity of the activity and ensuing fall.
219 Familiarity with the classification is likely to improve this. In addition the classification was
220 unable to consider medication state or the role of freezing and festination in falls.

221

222 We advocate clinical and research use of the FRAC. A clinical understanding of pre-fall
223 activity may help identify more specific and individualised fall prevention strategies. For
224 example, identification of an individual who falls almost exclusively during postural
225 transitions may mean therapy can be targeted towards function-based strengthening and
226 basic balance training. By contrast, if falls are categorised as combined, then it is possible
227 the individual may benefit from gait re-education, higher balance training, dual task training
228 or cueing strategies to ameliorate the gait dysfunction which is possibly a contributing
229 factor.

230

231 In addition, the FRAC may be used to provide a formal approach to fall classification which
232 can be charted over time, but also, most importantly for early identification of falls with the
233 intention of preventing the slide from an incidental 'high level' (advanced) fall to clinically
234 significant and concerning 'postural transition' (transitional) fall. This common clinical
235 assumption; that a high level fall may be a precursor to future lower level transitional falls,
236 needs to be tested, and the FRAC is a more appropriate means to investigate this theory
237 than classification by incidence.

238

239 Results from this study confirm the view that fallers vary phenotypically which is not evident
240 when using falls frequency as a classification. Although recurrent falls indicate greater
241 discrimination for pathology when compared to single falls [8], frequency as an outcome
242 alone is limiting. Mactier [5] found a significant association between ambulatory activity and
243 disease severity in incident PD was demonstrated when falls were categorised using the
244 FRAC, whereas the conventional classification, frequency, did not yield these findings [5].
245 This lends weight to the notion that falls (and fallers) are not homogeneous. Inspection of
246 the falls diaries revealed that participants were involved in a broad spectrum of physical
247 activities (not reported here) which ranged from climbing down a moss covered peat bog
248 before crossing a stream to simply getting out of bed. On the whole, transitional fallers
249 were depicted as being frail and more advanced in disease severity versus advanced fallers
250 who were portrayed as being more impulsive or lacking in insight. This has wider
251 implications, and highlights the need for future work to understand the different risk factors
252 conferred by grouping fallers and a potentially different approach to fall management
253 strategies. Nevertheless, falls researchers and clinicians almost exclusively measure falls by
254 frequency. This consistent approach enables between-study comparison and data
255 consilience, but comes at a cost of reducing meaningful interpretation. This ultimately limits
256 the advancement of falls research.

257

258 Strengths of this study include the design and methodology. Standardisation of the rating
259 process prevented bias. Use of 23 raters is uncommon in reliability studies and the variety of
260 professional backgrounds was an advantage. Limitations are that we did not measure test-
261 retest reliability or examine the accuracy of the patient's description of falls. It could be of

262 interest to correlate this classification with not only information gathered from more
263 objective measures such as accelerometers, but other classifications such as the postural
264 stability measures of the UPDRS [29]; all of which could be the focus of future work.

265

266 **CONCLUSIONS**

267 The FRAC can be reliably used to classify Parkinson's disease fallers. Classifying falls in this
268 way may discriminate between phenotypically different fallers which may increase the
269 likelihood of detecting, and subsequently strengthen risk factors for falls. Additionally, the
270 differences in fall types and outcomes may have implications for, and be used to guide fall
271 prevention strategies.

272

273 **Acknowledgement**

274 This research is supported by the National Institute for Health Research (NIHR) Newcastle
275 Biomedical Research Unit based at Newcastle upon Tyne Hospitals NHS Foundation Trust
276 and Newcastle University. The research was also supported by NIHR Newcastle CRF
277 Infrastructure funding. The views expressed are those of the authors and not necessarily
278 those of the NHS, the NIHR or the Department of Health. We would like to acknowledge
279 Dadirayi Mhiripiri and Rosie Morris for their assistance with data collection and Dr Brook
280 Galna for assistance with statistics.

281

282 **REFERENCES**

283 [1] Allen, N.E., Schwarzel, A.K. and Canning, C.G. (2013) 'Recurrent falls in Parkinson's
284 disease: a systematic review', *Parkinsons Dis*, p. 906274.

285 [2] Bloem, B.R., Grimbergen, Y.A.M., Cramer, M., Willemsen, M. and Zwinderman, A.H.
286 (2001) 'Prospective assessment of falls in Parkinson's disease', *J Neurol*, 248: 950-958.

287 [3] Kerr, G., Worringham, C., Cole, M., Lacherez, P., Wood, J. and Silburn, P. (2010)
288 'Predictors of future falls in Parkinson disease', *Neurology*, 75: 116-124.

289 [4] Pickering, R.M., Grimbergen, Y.A., Rigney, U., Ashburn, A., Mazibrada, G., Wood, B., Gray,
290 P., Kerr, G. and Bloem, B.R. (2007) 'A meta-analysis of six prospective studies of falling in
291 Parkinson's disease', *Mov Disord*, 22(13): 1892-900.

292 [5] Mactier, K., Lord, S., Godfrey, A., Burn, D. and Rochester, L. (2015) 'The relationship
293 between real world ambulatory activity and falls in incident Parkinson's disease: Influence of
294 classification scheme', *Parkinsonism and Related Disorders*. 21(3):236-42.

295 [6] Fasano, A., Plotnik, M., Bove, F. and Berardelli, A. (2012) 'The neurobiology of falls',
296 *Neurol Sci*, 33(6): 1215-23.

297 [7] Friedman, S.M, Munoz, B., West, S., Rubin, G.S. and Fried, L.P. (2002) 'Falls and Fear of
298 Falling: Which Comes First? A Longitudinal Prediction Model Suggests Strategies for Primary
299 and Secondary Prevention', *J Am Geriatr Soc*, 50:1329–1335.

300 [8] Mak, M.K. and Pang, M.Y. (2010) 'Parkinsonian single fallers versus recurrent fallers:
301 different fall characteristics and clinical features', *J Neurol*, 257(9): 1543-51.

302 [9] Canning, C.G., Sherrington, C., Lord, S.R., Close, J.C et al. (2015) 'Exercise for falls
303 prevention in Parkinson disease: A randomized controlled trial' , *Neurology*, 84(3): 304-12.

304 [10] Paul, S.S., Canning, C.G., Song, J., Fung, V.S., Sherrington, C. (2014) 'Leg muscle power is
305 enhanced by training in people with Parkinson's disease: a randomized controlled trial',
306 *Clinical Rehabilitation*, 28(3): 275–288.

- 307 [11] Ashburn, A., Stack, E.L., Ballinger, C., Fazakarley, L. and Fitton, C. (2008) ' The
308 circumstances of falls among people with Parkinson's disease and the use of Falls Diaries to
309 facilitate reporting', *Disability and Rehabilitation*, 30(16): 1205-1212.
- 310 [12] Nevitt, M.C., Cummings, S.R. and Hudes, E.S. (1991) 'Risk Factors for Injurious Falls: A
311 Prospective Study'. *Journal of Gerontology*. 46 (5): M164-170.
- 312 [13] Kelsey, J.L., Berry, S.D., Procter-Gray, E., Quach, L., Nguyen, U.D., Li, W., Kiel, D., Lipsitz,
313 L. and Hannan, M.T. (2010). 'Indoor and Outdoor Falls in Older Adults are Different: The
314 MOBILIZE Boston Study'. *J Am Geriatr Soc*, 58(11): 2135–2141.
- 315 [14] Bath, P. and Morgan, K. (1999). 'Differential Risk Factor Profiles for Indoor and Outdoor
316 Falls in Older People Living at Home in Nottingham, UK' *European Journal of Epidemiology*,
317 15 (1): 65-73.
- 318 [15] Blake, A.J., Morgan, K., Bendall, M., Dallosso, H., Ebrahim, S., Arie, T., Fentem, P. and
319 Bassey, E. (1988) 'Falls by Elderly People at Home: Prevalence and Associated Factors', *Age
320 and Ageing*, 17: 365-372.
- 321 [16] Campbell, J.A., Borrie, M. and Spears, G. (1989) 'Risk Factors for Falls in a Community-
322 Based Prospective Study of People 70 Years and Older', *Journal of Gerontology*, 44(4) :
323 M112-117.
- 324 [17] Lach, H. (1991) 'Falls in the elderly- reliability of a classification system', *Journal of the
325 American Geriatrics Society*, 39(2): 197-202.
- 326 [18] Waller, J.A. (1978) 'Falls among the Elderly- Human and Environmental Factors', *Accid.
327 Anat. & Prev*, 10: 21-33.


- 328 [19] Maki, B.E., Holliday, P.J. and Topper, A.K. (1994) 'Prospective Study of Postural Balance
329 and Risk of Falling in an Ambulatory and Independent Elderly Population', *Journal of*
330 *Gerontology*, 49(2):M72-M84.
- 331 [20] Wild, D., Nayak, U. and Isaacs, B. (1981) 'Description, Classification and Prevention of
332 Falls in Old People at Home', *Rheumatology and Rehabilitation*, 20: 153-159.
- 333 [21] Kelsey, J.L., Procter-Gray, E., Berry, S.D., Hannan, M.T., Kiel, D., Lipsitz, L. and Li, W.
334 (2012) 'Re-evaluating the Implications of Recurrent Falls in Older Adults: Location Changes
335 the Inference'. *J Am Geriatr Soc.* 60(3): 517–524.
- 336 [22] Khoo, T.K, Yarnall, A.J, Duncan, G.W., Coleman, S., O'Brien, J.T., Brooks, D.J, et al. (2013)
337 'The spectrum of non-motor symptoms in early Parkinson disease', *Neurology*, 80 (3): 276-
338 281.
- 339 [23] Hughes, A.J., Daniel, S.E., Kilford, L. and Lees, A J. (1992) 'Accuracy of clinical diagnosis
340 of idiopathic Parkinson's disease: a clinico-pathological study of 100 cases', *Journal of*
341 *Neurology, Neurosurg Psychiatry*, 55(3):181-4.
- 342 [24] Davalos-Bichara, M., Lin, F.R., Carey, J.P., Walston, J.D., Fairman, J.E., Schubert, M.C.,
343 Barron, J.S., Hughes, J., Millar, J.L., Spar, A., Weber, K.L., Ying, H.S., Zackowski, K.M., Zee,
344 D.S. and Agrawal, Y. (2013) 'Development and validation of a falls-grading scale', *J Geriatr*
345 *Phys Ther*, 36(2): 63-7.
- 346 [25] Light R.J. (1971) 'Measures of response agreement for qualitative data: Some
347 generalizations and alternatives'. *Psychological Bulletin*, 76(5):365–377.
- 348 [26] Hallgren, K. (2012) 'Computing Inter-Rater Reliability for Observational Data: An
349 Overview and Tutorial', *Tutor Quant Methods Psychol*, 8(1):23-34.

350 [27] Petrie, A., Sabin, C. (2009) 'Medical Statistics at a Glance', 3rd ed. Wiley-Blackwell.

351 [28] Mackenzie, L., Byles, J. and Higginbotham, N. (2002) 'Reliability of the Home Falls and
352 Accidents Screening Tool (HOME FAST) for identifying older people at increased risk of falls'
353 Disability and Rehab, 24 (5): 266-274.

354 [29] Goetz, C.G., Tilley, B.C., Shaftman, S. R. et al. (2008) 'Movement Disorder Society-
355 Sponsored Revision of the Unified Parkinson's Disease Rating Scale (MDS-UPDRS): Scale
356 Presentation and Clinimetric Testing Results'. Movement Disorder Society, 23 (15): 2129-
357 2170.

358



359
360 Figure 1: Falls Classification Continuum

361
362
363
364

365 Table 1: Fall-Related Activity Classification (FRAC) category descriptors

1. Advanced	2. Combined	3. Transitional
Involves a complex, high risk motor activity	Involves everyday walking activities including stair climbing or combined movements	Involves a basic transition from one posture to another
There is a significant environmental challenge that would explain the fall	Combined movements include moving from one position to another in a more challenging environment	Simple, safe and easy tasks
Unfamiliar indoor/outdoor environment		
e.g. Skiing, hill walking, slipping on ice	e.g. working in the garden, turning whilst walking, carrying heavy objects	e.g. rising from a chair, sitting on a sofa
The fall is expected in an age-matched non-PD person	The fall is not expected in age-matched non-PD person; it is a result of underlying physiological deficits	The fall is not expected in age-matched non-PD person

366
367
368
369
370

Table 2: Scenarios resulting in poor inter-rater reliability

Contentious falls scenarios			% agreement			Potential causes of disagreement based on category descriptors (<i>in italic</i>)		
Where were you?	What were you doing?	What do you think caused the fall?	1	2	3	Category 1	Category 2	Category 3
Walking the dog in the woods at the park	Trying to walk and manage lead/ bag and walking stick	Tripped over a tree root while not paying attention to what I was doing. Ground was muddy + slippy after the thaw	57	43	0	The woods at the park are an unfamiliar environment. Managing the lead/bag, tripping over tree root, slippy mud are all significant environmental challenges that would explain the fall	Walking the dog is a <i>simple everyday walking activity</i> . You would <i>not expect the fall in age-matched non-PD person</i>	
In the back lane	Mounting bike	Lost balance	39	61	0	Mounting a bike is a complex, high risk motor activity	Mounting a bike is a <i>combined movement, including moving from one position to another</i> The fall is <i>not expected in age-matched non-PD person</i>	
Fall on way to bed	I turned badly - overbalanced	Balance is very poor at times	0	57	43		Walking to bed is an everyday walking activity	Walking to bed is a <i>simple, safe and easy task</i> It involves a <i>basic transition from one posture to another</i>
Kitchen	Trying to turn around to go out of the kitchen	Loss of balance	0	61	39		Trying to turn around to go out of the kitchen is an everyday walking activity	Walking in the kitchen is a <i>simple, safe and easy task</i> It involves a <i>basic transition from one posture to another</i>
In kitchen	Open fridge door	Tripping over dog	22	52	26	Tripping over the dog is a <i>significant environmental challenge that would explain the fall</i>	Opening the fridge door in the kitchen is an everyday walking activity and a combined movement	Opening fridge door is a <i>simple, safe and easy task</i>

Gold standard in bold (classified by AR)

Table 3: Kappa values

Clinical Subgroup	N	Light's Kappa (CI_{2.5}, CI_{97.5})
All	23	.807 (.732, .870)
Physiotherapists	11	.822 (.750, .885)
Physicians	7	.821 (.747, .880)
Researchers	5	.780 (.697, .863)
Clinicians (physios & medics)	18	.815 (.744, .872)
Regular contact with fallers	16	.816 (.739, .878)
Falls experts	6	.810 (.697, .882)