

Newcastle University e-prints

Date deposited: 17th June 2010

Version of file: Author final

Peer Review Status: Peer Reviewed

Citation for published item:

Fowler HJ, Kilsby CG. [A regional frequency analysis of United Kingdom extreme rainfall from 1961 to 2000](#). *International Journal of Climatology* 2003,**23** 11 1313-1334.

Further information on publisher website:

<http://www.agu.org/> (*Website*)

Publishers copyright statement:

This paper was originally published by the American Geophysical Union, 2003 .

Use Policy:

The full-text may be used and/or reproduced and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not for profit purposes provided that:

- A full bibliographic reference is made to the original source
- A link is made to the metadata record in Newcastle E-prints
- The full text is not changed in any way.

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

<p>Robinson Library, University of Newcastle upon Tyne, Newcastle upon Tyne. NE1 7RU. Tel. 0191 222 6000</p>

A regional frequency analysis of United Kingdom extreme rainfall from 1961 to 2000

Fowler, H.J.* and Kilsby, C.G.

Water Resource Systems Research Laboratory, School of Civil Engineering and Geosciences,
University of Newcastle upon Tyne, UK

Dr Hayley J Fowler
WRSRL
School of Civil Engineering and Geosciences
Cassie Building
University of Newcastle
Newcastle upon Tyne
NE1 7RU

Tel: 0191 222 8599
Fax: 0191 222 6669
h.j.fowler@ncl.ac.uk

Sponsor

SWURVE (Sustainable Water: Uncertainty, Risk and Vulnerability Estimation in Europe) project,
funded under the EU Environment and Sustainable Development programme, grant number
EVK1-2000-00075

Abstract

Multi-day rainfall events are an important cause of recent severe flooding in the UK and any change in the magnitude of such events may have severe impacts upon urban structures such as dams, urban drainage systems and flood defences and cause failures to occur. Regional pooling of 1-, 2-, 5- and 10-day annual maxima for 1961 to 2000 from 204 sites across the UK is used in a standard regional frequency analysis to produce GEV growth curves for long return-period rainfall events for each of nine defined climatological regions. Temporal changes in 1-, 2-, 5- and 10-day annual maxima are examined with L-moments using both a 10-year moving window and fixed decades from 1961-70, 1971-80, 1981-90 and 1991-2000. A bootstrap technique is then used to assess uncertainty in the fitted decadal growth curves and to identify significant trends in both distribution parameters and quantile estimates.

There has been a two-part change in extreme rainfall event occurrence across the UK from 1961-2000. Little change is observed at 1- and 2-day duration, but significant decadal level changes are seen in 5- and 10-day events in many regions. In the south of the UK, growth curves have flattened and 5- and 10-day annual maxima have decreased during the 1990s. However, in the north, the 10-day growth curve has steepened and annual maxima have risen during the 1990s. This is particularly evident in Scotland. The 50-year event in Scotland during 1961-1990 has become an 8-, 11- and 25-year event in the Eastern, Southern and Northern Scotland pooling regions respectively during the 1990s. In northern England the average recurrence interval has also halved. This may have severe implications for design and planning practices in flood control.

Keywords

Region, Technique(s), Climate variable(s), Rainfall, Extremes, Flooding, Climate change, Recurrence interval

Introduction

The autumn and winter of 2000/2001 provided the wettest months on record in the UK (Marsh, 2001) and widespread flooding throughout the last decade has enhanced the public concern that such heavy and prolonged rainfall events are a result of global warming. Analyses of rainfall trends in observed data (e.g. Bradley *et al.*, 1987, Diaz *et al.*, 1989, Groisman *et al.*, 1999) suggest a trend towards increased rainfall and enhanced variability in high latitudes of the northern Hemisphere (e.g. Easterling *et al.*, 2000), particularly in winter. Significant positive trends in intensity have been observed in the UK (Osborn *et al.*, 2000; Lamb, 2001), Europe (Brunetti *et al.*, 2000; Frei and Schar, 2001) and worldwide (Karl and Knight, 1998; Iwashima and Yamamoto, 1993; Zhai *et al.*, 1999). These observations show agreement with the results of climate model integrations which predict increases in both the frequency and intensity of heavy rainfall in the high latitudes under enhanced greenhouse conditions (e.g. Murphy and Mitchell, 1995, McGuffie *et al.*, 1999). The amplification of the global water cycle through global warming may have severe consequences in the UK, particularly in terms of increases in significant flood events.

This study considers changes in extreme rainfall event frequency and intensity across the UK. An analysis of observed trends from 1961-1995 by Osborn *et al.* (2000) indicate increases in the number of heavy rainfall days along with increases in the rainfall amount on those days, particularly in winter months. Additional analyses at 3 UK case-study locations by Lamb (2001) suggest that the frequency of longer duration extreme rainfalls has been increasing since the 1960s. These trends are supported by analysis of General Circulation Model (GCM) simulations and Regional Climate Model (RCM) integrations. Palmer and Raisanen (2002) analysed 19 GCM simulations, and estimated that the probability of winter rainfall exceeding two standard deviations above normal will increase by factors of five and three respectively over northern and southern parts of the UK by 2100. RCM integrations over the period from 2080-2100 (Jones

and Reid, 2001) also suggest that there will be dramatic increases in the heaviest rainfall events over Britain.

The observed increase in extreme rainfall has already caused severe impacts in both the UK and northern Europe, including major fluvial flooding (Marsh, 2001; Lamb, 2001) and landslides affecting rail networks, notably the East Coast line at Doncaster and fatal incidents in Scotland and northern England. A characteristic of these events has been their multi-day nature, with unremarkable one-day totals (Lamb, 2001). These changes to the spatial and temporal distribution of extreme rainfall events in the UK are very important in the design of urban structures such as drainage systems, dams, spillways and flood control measures, for example, and may cause increased failure of such systems. Therefore, a re-evaluation of design estimates is needed.

In general, two approaches have been taken to assess rainfall extremes. The first approach uses a percentile or quantile method to assess extreme rainfall (e.g. Karl and Knight, 1998; Osborn *et al.* 2000). In this approach daily rainfall records are sorted and classes defined to contain a certain percentage of the total number of rainfall events for a season or month. Each of the classes contains an equal amount of total rainfall and can therefore be thought of as amount quantiles. The second approach uses statistical distributions to define extremes with given return periods on an annual basis (e.g. Hennesey *et al.*, 1997; McGuffie *et al.*, 1999). In this method, estimation of the magnitude of long return-period rainfall events involves fitting an extreme value distribution to the AM series. This method produces return period estimates that are easily understood and can be used readily for design purposes.

In this study, a regional frequency analysis approach based on L-moments (Hosking and Wallis, 1997) is used to produce rainfall growth curves with an extreme value distribution. This involves the regional pooling of annual maxima (AM) to allow the estimation of long return-

period rainfall events when individual records are too short to allow their reliable estimation. Daily rainfall records from 204 sites across the UK for 1961-2000 are used to produce rainfall growth curves for 1-, 2- 5- and 10-day AM series for each of the nine homogeneous rainfall regions defined by Wigley *et al.* (1984).

This method of pooling is also used to examine temporal variability in L-moment ratios and AM series, utilising both a 10-year moving window and fixed decades from 1961-1970, 1971-1980, 1981-1990, and 1991-2000. This provides a comprehensive analysis of spatial and temporal changes in extreme rainfall event occurrence across the UK from 1961-2000.

Observed rainfall data

The study has been limited by necessity to daily data, as sub-daily data are not generally available with sufficient coverage and length of record. However, daily data are adequate for the purposes of this study, since attention is focussed on multi-day events.

Daily data for 110 stations (after Osborn *et al.*, 2000) from 1961-1995 were updated using data from the British Atmospheric Data Centre (<http://www.badc.rl.ac.uk/>) to give complete records from 1961-2000. Additional sites were then chosen such that each of the nine regions defined by Wigley *et al.* (1984) in their study of UK rainfall contained at least 20 stations. These regions are defined in Figure 1 and were originally defined to take account of physiographic character and spatially coherent rainfall variability, based on an average of seven sites per region. This gave a total of 204 stations across the UK chosen on the basis of record length and completeness, and to provide a good spatial coverage (see Figure 1).

Regionalisation

Analyses were performed using L-moments of the AM series to determine whether the regions of Wigley *et al.* (1984) are appropriate for an analysis of extreme rainfall. Firstly, the three L-moment ratios L-CV, L-skewness and L-kurtosis were determined for the AM series at each site using a routine from Hosking (1997). Figure 2 shows a plot of L-CV: L-skewness of the mean values for each of the nine pooling regions. It can be seen that generally, eastern regions display a greater L-CV value than western regions, suggesting higher variability in these regions. The highest L-skewness values are found in Southwest and Southeast England. These fall as a move is made northwards, to much lower values in North Scotland. In simple terms, this suggests that more intense rainfall events are experienced in southern regions of the UK. This is in line with the accepted variation in rainfall growth curves as expressed in the regions of the Flood Studies Report (1975).

A discordancy analysis was then performed to establish if the distributions of site AM series within each region were acceptably similar. The discordancy measure, D_i , (Hosking and Wallis, 1997) compares the L-moment ratios of a site with those of the pooling group as a whole, hence identifying sites with L-moment ratios that are unusual relative to the pooling group. A high value of the discordancy measure indicates that a site may be discordant within the pooling group, but this may be caused by only a few unusual rainfall events. D_i is formally defined by Robson and Reed (1999): if M is the number of sites in the pooling group and \mathbf{u}_i is a vector of the L-moment ratios at site i , then (1):

$$\mathbf{u}_i = (t_1, t_2, t_3)^T \quad (1)$$

where t_1 is L-CV, t_2 is L-skewness, t_3 is L-kurtosis and superscript T denotes the transpose of a vector. Thus defining two matrices \mathbf{U} (2) and \mathbf{A} (3) as:

$$\mathbf{U} = \frac{1}{M} \sum_{i=1}^M \mathbf{u}_i \quad (2)$$

$$\mathbf{A} = \sum_{i=1}^M (\mathbf{u}_i - \mathbf{U})(\mathbf{u}_i - \mathbf{U})^T \quad (3)$$

Then the discordancy measure D_i for site i is given by (4):

$$D_i = \frac{1}{3} M (\mathbf{u}_i - \mathbf{U})^T \mathbf{A}^{-1} (\mathbf{u}_i - \mathbf{U}) \quad (4)$$

where \mathbf{A}^{-1} is the inverse of matrix \mathbf{A} .

Critical values of the discordancy measure for each site in a pooling group based on a 10% significance level are suggested by Hosking and Wallis (1997). For a pooling group with more than 15 member sites, values of D_i higher than 3.0 show possible discordancy. Values of D_i greater than the critical value of 3 were only found for three sites. These were in the pooling regions of Northeast and Northwest England.

In the NEE pooling region the two discordant sites, Lockwood Reservoir and Whitby Coastguard, have D_i values of 3.29 and 3.15 respectively. At Lockwood Reservoir the high discordancy measure arises due to a single, unusually high rainfall event in 1976 when 104.6 mm fell in 24-hrs, and similarly at Whitby Coastguard. In the NWE pooling region, the site at Appleby is group discordant with a D_i of 3.06. At this site, again, the high discordancy measure results from a single, unusually high, AM. In this case, a heavy rainfall event on the 17th July 1983 produced 97.8 mm rainfall in 24-hrs. In each case, a single heavy rainfall event

was found to have caused unusually high values of L-kurtosis, and so no adjustment to the pooling groups was considered necessary.

Regional Frequency Analysis

Methodology

A regional frequency analysis approach based on L-moment methods (Hosking and Wallis, 1997) was taken to generate rainfall growth curves for 1-, 2-, 5- and 10-day events for each of the 9 regions. For each site, annual rainfall maxima for 1-, 2-, 5- and 10-day events are standardised using the median event (*RMED*), calculated using data from the period 1961-1990. This removes site-specific factors from the regional analysis and is the same variable as that used in the FORGEX (Focused Rainfall Growth Extension) method (Reed *et al.*, 1999) of the Flood Estimation Handbook (FEH) (IH, 1999). L-moment ratios derived from single site analyses within a region are then combined by regional averaging, weighted according to record length (see Hosking and Wallis, 1997). Thus, giving an example formula for L-CV (5):

$$LCV_{pooled} = \sum_{i=1}^N w_i LCV_i \quad (5)$$

where N is the number of sites in the pooling group and the weight w_i is an effective record length at the i^{th} site defined by (6):

$$w_i = \frac{n_i}{\sum_{i=1}^N n_i} \quad (6)$$

The denominator is the total number of station-years of record in the pooling group, while the numerator is the number of station-years at the i^{th} site. The weighted average L-Skewness and L-Kurtosis moment ratios are derived in the same way. The station-year approach to estimating long return-period events is avoided as this assumes that site records are mutually independent; only valid for very widely scattered sites and thus unjustified in this application.

The classical L-moments approach is then used to fit the GEV (Generalised Extreme Value) distribution for each AM series by matching the sample L-moments to the distribution L-moments. The GEV distribution is widely used in extreme event frequency analysis in the UK (e.g. FEH) and is used in preference to the Gumbel distribution as the literature increasingly suggests that the distribution of extreme events may be heavier tailed.

The changing shape of the standardised AM series as it moves from 1-day to 10-day rainfall totals and a comparison of the shape of the unfitted distributions between different regional pooling groups can be seen in Figure 3. Although most of the growth curves approximate a straight line, and therefore the Gumbel distribution, others have significant curvature. This is particularly apparent in the SEE region, where the 1-, 5- and 10-day AM series are significantly curved, and justifies the choice of the GEV distribution in fitting. The curvature is also evident to a lesser extent in the 10-day AM series of both the SWE and NS pooling regions and is possibly due to the existence of two mechanisms of rainfall; frontal (flat) and convective (curved upper section). It can be observed that, for 10-day totals in particular, many of the AM distributions exhibit significant curvature. This is especially prevalent in the south of England and may be important in the prediction of future rainfall extremes.

The GEV distribution has three parameters and is described by (7):

$$x(F) = \xi + \frac{\alpha}{k} \{1 - (-\ln F)^k\} \quad (k \neq 0) \quad (7)$$

where ξ is the location parameter, α the scale parameter, k the shape parameter and F refers to a given quantile.

A growth curve was fitted for each AM series using the regionally averaged site L-moment ratios. The fitted growth curve is given by (8):

$$x(F) = 1 + \frac{\beta}{k} \{(\ln 2)^k - (-\ln F)^k\}$$

where

$$\beta = \frac{\alpha}{\xi + \frac{\alpha}{k} \{1 - (\ln 2)^k\}} \quad (8)$$

The parameter k is estimated from the L-skewness (Hosking *et al.*, 1985) (9):

$$k \approx 7.8590c + 2.9554c^2$$

where

$$c = \frac{2}{3 + t_3} - \frac{\ln 2}{\ln 3} \quad (9)$$

The parameter β is estimated using L-CV (Hosking and Wallis, 1997) as (10):

$$\beta = \frac{kt_2}{t_2 \{ \Gamma(1+k) - (\ln 2)^k \} + \Gamma(1+k)(1-2^{-k})} \quad (10)$$

where Γ denotes the gamma function, t_2 is the L-CV L-moment ratio and t_3 is the L-Skewness L-moment ratio.

The fitted distribution plotting positions on a variate versus Gumbel reduced variate plot are determined according to the Gringorten (1963) formulae (11):

$$F_i = \frac{i + 0.44}{N + 0.12} \quad (11)$$

where F_i is the non-exceedance probability, i the rank in increasing order, and N the number of annual maxima in the pool. The Gumbel reduced variate is defined by (12):

$$y = -\ln(-\ln F_i) \quad (12)$$

The fitted growth curves are shown in Figure 4. Figure 5 gives an approximation of the pooled AM rainfall in millimetres by using the geometric mean *RMED* of all stations in the pooling group for a particular event duration during the period 1961-1990 as a multiplier in each case. Fitted GEV distribution parameters for the nine regions can be found in Table 1.

Estimating uncertainty

An estimate of uncertainty in return period predictions of the pooled growth curve gives some confidence in the use of the growth curve for design purposes. Two methods have been used in this analysis. The first, the pooled uncertainty measure, *PUM* (Robson and Reed, 1999),

addresses the variation in growth curves within a region. The *PUM* is a weighted average of the differences between each site growth factor and the pooled growth factor measured on a logarithmic scale. The pooled uncertainty measure for return period T , PUM_T , is defined by (13) (Robson and Reed, 1999):

$$PUM_T = \sqrt{\frac{\sum_{i=1}^N n_i (\ln x_{T_i} - \ln x^P_{T_i})^2}{\sum_{i=1}^N n_i}} \quad (13)$$

where N is the number of sites in the pool, n_i is the record length in years of the i^{th} site, x_{T_i} is the T -year growth factor for the i^{th} site, and $x^P_{T_i}$ is the T -year pooled growth factor for the i^{th} site.

The PUM_T for $T = 2, 5, 10, 25$, and 50 can be found in Table 2, reflecting the average uncertainty in the T -year growth factor at each return period. An estimate of the regional average uncertainty in millimetres is given by multiplying by the regional mean *RMED* in each case. The uncertainties are small, suggesting coherent pools, with typical values of *PUM* ranging from 6 mm for 1-day 50 year events up to 12 mm for 10-day 50 year events. The design storm can thus be estimated for 1-, 2-, 5- and 10-day rainfall totals for any part of the UK by using the correct regional growth curve and multiplying by the station *RMED*. To allow for uncertainty in the pooled growth curve, the appropriate PUM_T should be multiplied by the station *RMED* and then added to the previous design storm estimate.

The second measure allows for the effect of spatial dependence, where single major storm events give rise to annual maxima at several sites in the same region. A bootstrap method

(Efron, 1979) was used where one year of record at a time was removed from the pooled regional data and the growth curve refitted for each decade. This gave uncertainty limits around the growth curves for each region which are larger than those derived using the PUM. The combined uncertainty envelope is then plotted for each growth curve.

Estimating temporal change in extreme rainfall

Methodology

Changes in the regional growth curve parameters and L-moment ratios during the period from 1961-2000 were examined using a 10-year moving window. For each of the 204 sites, annual maxima are found for 1-, 2-, 5- and 10-day rainfall totals. If the i th AM rainfall amount at site j during decade k is denoted by P_{ijk} then the standardized annual maxima are defined as (after Reed *et al.*, 1999) (14):

$$X_{ijk} = \frac{P_{ijk}}{RMED_{jk}} \quad (14)$$

where $RMED_{jk}$ denotes the median of annual maxima at site j during decade k . For each decade in a moving 10-year window from 1961 to 2000, the L-moment ratios were determined using a routine from Hosking (1997) and examined for trends.

Further analysis was undertaken by splitting the records into four separate decades 1961-1970, 1971-1980, 1981-1990 and 1991-2000 and standardising by $RMED$ as before. For each region, the standardized AM data are then pooled for each separate decade and GEV distributions fitted. Finally, each AM is multiplied by the mean $RMED$ of all stations in that pool for that decade, i.e. (15):

$$\hat{P}_{ijk} = X_{ijk} \sum_{j=1}^{j=n} RMED_{jk} / n \quad (15)$$

where \hat{P}_{ijk} gives an approximation of the pooled AM rainfall in millimetres.

Change in L-moment ratios

The L-moment ratios are a direct measure of the AM distribution and, as such, provide a better illustration of changes than fitted GEV parameters themselves. Figure 6 shows variation and trend in L-moment ratios for regions and durations where the change is the most significant. Results indicate that firstly, at 1- and 2-day durations, there has been a shift in some western regions to lower L-moment ratios from 1961 to 2000. This is especially prevalent in the SWE and NI pooling regions (Figure 6; (a, b)). However, this change is not seen in other western regions such as NWE and NS. Secondly, at 5- and 10-day durations, in particular 10-days, there is a two-part shift across the UK. In northern and western regions (NS, SS, ES, NWE and NI) there has been a move to higher L-moment ratios, particularly in ES (see Figure 6(d)). In southern and eastern regions (CEE, NEE, SEE and SWE), L-moment ratios have been declining throughout the period and especially in the most recent decade (see Figure 6(c)).

In simple terms this represents a change in extreme rainfall properties, with increased variability (rising L-CV) and increased intensity (rising L-skewness) of 5- and 10-day rainfall events in northern and western regions, and the opposite change occurring in southern and eastern parts of the UK. A similar shift in western regions indicates decreased variability in the 1-day rainfall event.

Change in growth curve shape

The shape of the fitted GEV distribution has changed over the last 40 years in some regions. These changes are markedly different between the south and north of the UK and are mainly seen at 5- and 10-day duration.

In the SEE, SWE and CEE pooling regions the growth curve has become flatter over the last decade from 1991 to 2000. This can be seen in the 5- and 10-day rainfall annual maxima, particularly in the SEE pooling region. Growth curves for 10-day duration in SEE and SWE are shown in Figures 7(a) and 7(b) respectively, along with their uncertainty estimates. These overlap, hence the change cannot be considered significant. In the NEE and NWE pooling regions the opposite change occurs with the growth curve increasing in gradient for 5- and 10-day rainfall annual maxima. Growth curves for the 10-day duration are shown in Figures 7(c) and 7(d) respectively. Uncertainty limits show that the changes are not significant.

In Scotland, dramatic changes in the shape of the growth curve occur in the east. In ES (Figure 8) the growth curve steepens markedly at all durations and for all decades. These changes are significant as the uncertainty limits are clearly separated from the growth curves in previous decades, particularly for the 5-day duration (significant for all return period estimates) and the 10-day duration (significant above the 10-year return period estimate). In the North and South Scotland pooling regions (not shown), however, there has been little change in the shape of the distribution over the last 40 years.

Change in magnitude of annual maxima (*RMED*)

Table 3 gives mean *RMED* values for each pooling region at each duration and for each decade. It can be seen that in general, and particularly in the north and west of the UK, the 1980s and 1990s have seen a changing *RMED*, with a shift upwards in most cases. This is especially obvious for 5- and 10-day rainfall events, where 1991-2000 shows the highest mean *RMED* for a 10-day event in every region excepting SEE, suggesting that it is at longer durations that change is occurring. This is most prominent in Scotland. In SS and ES the highest mean *RMED* values for all durations occur during the decade from 1991 to 2000.

In southern and central England recent events during the 1980s and 1990s have been less extreme than events during the 1960s and 1970s. The AM series for SWE, SEE and CEE reveal many extreme events in the 1960s and 1970s. The decade from 1961-1970 was in fact highly unusual, the period from 1961-1964 being extremely dry across the UK, but thereafter becoming extremely wet. These two extremes of dry and wet, found in the same decade, may account for the unusual shape of the growth curve found in both the regions of SWE and NI (Figures 9(a) and 9(b)) during the 1960s.

Unusually high annual maxima during the 1960s and 1970s can particularly be seen in the AM series for the SEE region (Figures 9(c) and 9(d)). The largest AM during the decade from 1961-1970 occur in 1968. In the 1960s, high 2-day AM and smaller 1-day AM are associated with the event of the 14th and 15th September 1968. During the 1970s, there are high annual maxima in 1973. These are related to a single event on the 20th September and provide very large 1-day rainfall totals at some sites. At Manston (Kent) for example, there is 160.8 mm rainfall, 5.6 times larger than the decadal site median. Rainfall approximating 100 mm also occurred at Boxley (Kent), 102.1 mm, Faversham (Kent), 99.1 mm, and Wye (Kent), 127.5 mm.

Although in southern and central England recent rainfall events have been less extreme than events during the 1960s and 1970s, in the rest of the nine regions defined by Wigley *et al.* (1984) a different picture emerges. In northern regions there has been a move towards increasing extremes in recent years. Nowhere is this more prominent than eastern Scotland.

In Eastern Scotland the growth curve during the 1990s provides a dramatic departure from the growth curve during the previous decades, and is enhanced considerably by the increase in *RMED* (see Figure 10). This is particularly apparent for the 5- and 10-day growth curves, due to a large increase in extreme rainfall events during the 1990s. This increase can be substantially explained by rainfall events during September 1995 and October 1993.

In September 1995, the north and east of Scotland were the wettest regions of the UK (181mm, 299 per cent of normal in east Scotland) (Cullum, 1996). On the 7th to the 8th September, 67 mm rainfall fell at Aberdeen. This was followed by a further 70 mm at Glenlivet on the 9th/10th September. A further depression tracked across the UK on the 10th to the 11th September and Kinloss received 69 mm in 24 hours on the 11th/12th September. Repeated heavy downpours in northeast Scotland caused serious flooding along the southern shore of the Moray Firth and around Aberdeen. During the first 11 days of September 272mm rain fell at Kinloss, for example, 13 times the normal for such a period. At Dyce the monthly total was 234 mm (344 percent of the 1961-90 average) (Weather Log, 1995).

In October 1993, there was widespread heavy rainfall and thunder over the first few days caused by a complex low centred over England (Weather Log, 1993). At Dyce, the total for the month was 119 mm (145 percent of the 1961-90 average) and at Leuchars the total was 112 mm (198 percent of the 1961-90 average) (Weather Log, 1993).

There has also been a recent increase in extreme event occurrence in the pooling regions of SS, NS and NI, particularly at 5- and 10-day durations. The 10-day growth curves multiplied by regional mean *RMED* for NS and SS are shown in Figures 11(a) and 11(b) respectively. Here, however, the increase starts as early as the 1980s and is due to an increase in the magnitude of events (i.e. increasing *RMED*) rather than changes to the shape of the growth curve.

This increase is also seen in NEE and NWE for 5- and 10-day AM series. Figures 11(c) and 11(d) show the 10-day return period estimates for an average site in the NEE and NWE regions respectively. In the 1990s there has been an increase in long duration, high magnitude events caused by both changing shape of the growth curve and increasing *RMED*.

Change in return period estimates

The above changes can be better seen in Table 4 which compares the 100-year event for 1961-1990 with that of 1991-2000 for 5- and 10-day durations. In southern regions, there is either little or negative change in the magnitude of the 100-year event. However, in northern pooling regions during the 1990s the 100-year event shows an increase in magnitude when compared to the 1961-1990 average. This change is very prominent in Eastern Scotland, with a 75% increase in rainfall amount for the 100-year event. This may substantiate the recent move in Scotland to use the 200-year event for design and planning purposes (Fleming, 2001).

Another, and the more common, way of looking at this type of change is as a change in recurrence interval. Table 5 shows the change in average recurrence interval for a 50-year event from for 5- and 10-day durations for the 1961-90 time period with the most recent decade from 1991 to 2000. It can be seen that the average recurrence interval has increased in southern pooling regions but decreased substantially in northern pooling regions. The 50-year event in Scotland during 1961-1990 has become an 8-year, 11-year and 25-year event in the ES, SS and NS pooling regions respectively during the 1990s. In northern England (NEE and NWE pooling regions) the average recurrence interval has also halved. This may have severe implications for design and planning practices for flood control.

Discussion and conclusions

This study has shown that the frequency of extreme rainfall has changed over parts of the UK in the period 1961-2000. This may be due to either or both natural climatic variability or climate change. However, there are several caveats to the approach taken in this research and these are discussed further here.

Firstly, although most procedures for rainfall frequency estimation are based on the analysis of AM series, this contradicts theoretical advice which invariably recommends peak-over-threshold or POT analysis. POT methods include all large events and exclude annual maxima that may be small and thus misleading in an extreme analysis. Unfortunately however, the treatment of missing data in the POT approach is demanding and determining which peaks to exclude during the same rainfall event can be very time consuming, so the AM series approach was adopted in the first instance.

Secondly, no explicit account has been taken in this analysis of the spatial dependence of rainfall events between stations in a given region. It is clear, however, that some regions are affected by a single storm event giving rise to large totals at several sites. Many researchers have examined the problem of spatial dependence in the prediction of rainfall extremes (e.g. Hosking and Wallis, 1988; Dales and Reed, 1989). Dales and Reed (1989) account for spatial dependence by reducing the 'network' (pooling group) to an effective number of independent sites based on the number of sites in the network, area of the network and the duration of rainfall extremes. This allows the growth curve to be shifted a fixed distance to the right to account for spatial dependence. However, this method has not been extensively validated. Hosking and Wallis (1988) used a Monte Carlo simulation to assess the effect of realistically specified inter-site dependence on growth curve estimation in a regional flood frequency analysis. They found that; (a) any bias in flood quantile estimates is unchanged by the presence of inter-site dependence; (b) regional heterogeneity exerts a stronger effect on the growth curve than inter-site dependence, and, moreover; (c) even when both heterogeneity and inter-site dependence are present, regional frequency analysis is more accurate than at site analysis. Therefore, as a definitive methodology to account for spatial dependence is unavailable, this research relies on a bootstrap to estimate the likely error.

Thirdly, the assumption of ‘quasi-stationarity’ has been made in the decadal analysis. The selection of a ten-year period represents an arbitrarily chosen compromise between assuming stationarity and acquiring a larger data set to allow longer return periods to be estimated.

These assumptions aside, the major conclusions of this study are:

1. There have been significant but regionally varying changes in extreme rainfall event occurrence across the UK in the last forty years;
2. Prolonged heavy rainfall events are increasing in northern and western regions. Growth curves have become steeper and annual maxima have increased during the 1990s due to both the growth curve changes and a large increase in regional mean *RMED*. This is particularly evident in eastern Scotland, where the recent decade from 1991-2000 provides a significant departure from previous return period estimates.
3. In the south (SEE, SWE and CEE) growth curves have become flatter and 5- and 10-day annual maxima have decreased during the 1990s, despite increasing *RMED* in most regions.

It has been shown that multi-day, prolonged heavy rainfall events are increasing in northern and western parts of the UK. These changes have implications for the design and maintenance of infrastructure such as urban drainage systems and flood control measures. Return period estimates presented here for different regions of the UK, using the most recent rainfall data, will allow the re-assessment of the risk of failure of existing structures and facilitate the design of new structures incorporating better risk or uncertainty estimates.

A remaining issue is the question of what period of record should be used to formulate design standards in a transient climate. Many researchers have questioned the validity of giving equal

weight to both historical and modern hydrometric data in return period estimation (e.g. Marsh, 1996) given the changing pattern and increasing persistence of heavy rainfall events. For example, current practice is to rely on a climatological normal period such as 1961-1990. FEH relies on historic data extending further back than 1961. There are strong indications from this work and elsewhere that changes have occurred in the 1980's and 1990's, which casts doubt on this approach. The alternatives are to use either a shorter period such as 1991-2000 which may be more representative of future conditions, or to accommodate underlying non-stationarity in some way. Both of these approaches are fraught with uncertainties, and further research is needed to establish a robust approach which avoids the assumption of stationarity whilst using sufficient records to obtain reliable estimates.

Future possibilities

It is anticipated that the research presented in this paper will be built upon to examine the further possibilities:

- Seasonal change in rainfall extremes (Fowler and Kilsby, 2003); this may be particularly important given the recent severe autumn flood events of 2000 and 2001 (Lamb, 2001), and notable changes in the seasonal distribution of rainfall quantiles shown by Osborn *et al.* (2000).
- Using peak-over-threshold data to produce revised growth curve estimates.
- Comparing the observed return period estimates with those generated using a regional climate model such as HadRM3 to develop a methodology that can produce accurate growth curves for the future climate given current information.
- Linking trends in rainfall extremes to trends in floods using various case-studies.

Acknowledgements

We thank Tim Osborn for the use of the daily rainfall data set and the British Atmospheric Data Centre (BADC) for the most recent rainfall data. This work is part of the SWURVE (Sustainable Water: Uncertainty, Risk and Vulnerability Estimation in Europe) project, funded under the EU Environment and Sustainable Development programme, grant number EVK1-2000-00075. We thank both Duncan Reed and Paul Bates for comments that helped to improve this paper.

References

1. Bradley RS, Diaz HF, Eischeid JK, Jones PD, Kelly PM, Goodess CM. 1987. Precipitation fluctuations over Northern Hemisphere land areas since the mid-19th century. *Science*. **237**: 171-275.
2. Brunetti, M., Buffoni, L., Maugeri, M. and Nanni, T. 2000. Precipitation intensity trends in northern Italy. *International Journal of Climatology*. **20**: 1017-1031.
3. Cullum, D. 1996. Some features of Northern Hemisphere weather during autumn 1995. *Weather*. **51**: 141-143.
4. Dales, M.Y. and Reed, D.W. 1989. Regional flood and storm hazard assessment. *Institute of Hydrology Rep. No. 102*. IH, Wallingford, UK.
5. Diaz HF, Bradley RS, Eischeid JK. 1989. Precipitation fluctuations over global land areas since the late 1800s. *Journal of Geophysical Research*. **94**: 1195-1240.
6. Easterling, D.R., Meehl, G.A., Parmesan, C., Changnon, S.A., Karl, T.R. and Mearns, L.O. 2000. Climate extremes: Observations, modeling, and impacts. *Science*. **289**: 2068-2074.
7. Efron, B. 1979. Bootstrap methods: another look at the jack-knife. *Annals of Statistics*. **7**: 1-26.
8. Fleming, G. 2001. Managing flood risk: past, present and future. *Flood Risk in a Changing Climate. Discussion Meeting at the Royal Society, London, 21st – 22nd November, 2001*.
9. Fowler, H.J. and Kilsby, C.G. 2003. Implications of changes in seasonal and annual extreme rainfall. *Geophysical Research Letters (in press)*.
10. Frei, C., and Schar, C. 2001. Detection probability of trends in rare events: Theory and application to heavy precipitation in the alpine region. *Journal of Climate*. **14**: 1568-1584.
11. Gringorten, I.I. 1963. A plotting rule for extreme probability paper. *Journal of Geophysical Research*. **68**: 813-814.
12. Groisman PY, Karl TR, Easterling DR, Knight RW, Jamason PF, Hennessy KJ, Suppiah R, Page CM, Wibig J, Fortuniak K, Razuvaev VN, Douglas A, Forland E, Zhai PM. 1999. Changes in the probability of heavy precipitation: Important indicators of climatic change. *Climatic Change*. **42**: 243-283.
13. Hennesey, K.J., Gregory, J.M. and Mitchell, J.F. 1997. Change in daily precipitation under enhanced greenhouse conditions. *Climate Dynamics*. **13**: 667-680.
14. Hosking, J.R.M. 1997. Fortran routines for use with the method of L-moments. *Research Report RC 20525 (90933) Version 3.02*. IBM Research Division: New York. 33pp.
15. Hosking, J.R.M. and Wallis, J.R. 1988. The effect of intersite dependence on regional flood frequency analysis. *Water Resources Research*. **24**: 588-600.
16. Hosking, J.R.M. and Wallis, J.R. 1997. *Regional frequency analysis: an approach based on L-moments*. Cambridge University Press. 224pp.
17. Hosking, J.R.M., Wallis, J.R. and Wood, E.F. 1985. Estimation of the generalised extreme-value distribution by the method of probability-weighted moments. *Technometrics*. **27**: 251-61.
18. IH 1999. *Flood Estimation Handbook*, 5 volumes, CEH Wallingford (formerly the Institute of Hydrology), Wallingford, UK.
19. Iwashima, T. and Yamamoto, R. 1993. A statistical analysis of the extreme events: long-term trend of heavy daily precipitation. *Journal of the Meteorological Society of Japan*. **71**: 637-640.

20. Jones, P.D. and Reid, P.A. 2001. Assessing future changes in extreme precipitation over Britain using Regional Climate Model integrations. *International Journal of Climatology*. **21**: 1337-1356.
21. Karl, T.R. and Knight, R.W. 1998. Secular trends of precipitation amount, frequency, and intensity in the United States. *Bulletin of the American Meteorological Society*. **79**: 231-241.
22. Lamb, R. 2001. To what extent can the October/November 200 floods be attributed to climate change. *DEFRA FD2304 Final Report*, 40pp.
23. Marsh, T.J. 1996. The 1995 drought - a signal of climatic instability? Proceedings of the Institution of Civil Engineers, Water, Maritime and Energy. **118**: 189-195.
24. Marsh, T.J. 2001. The 2000/2001 floods in the UK – a brief overview. *Weather*. **56**: 343-345.
25. McGuffie, K., Henderson-Sellers, A., Holbrook, N., Kothavala, Z., Balachova, O., Hoekstra, J. 1999. Assessing simulations of daily temperature and precipitation variability with global climate models for present and enhanced greenhouse climates. *International Journal of Climatology*. **19**: 1-26.
26. Murphy, J.M., and Mitchell, J.F.B. 1995. Transient response of the Hadley Centre coupled ocean-atmosphere model to increasing carbon-dioxide. Part II: Spatial and temporal structure of response. *Journal of Climate*. **8**: 57-80.
27. Osborn, T.J., Hulme, M., Jones, P.D. and Basnett, T.A. 2000. Observed trends in the daily intensity of United Kingdom precipitation. *International Journal of Climatology*. **20**: 347-364.
28. Palmer, T.N. and Raisanen, J. 2002. Quantifying the risk of extreme seasonal precipitation events in a changing climate. *Nature*. **415**: 512-514.
29. Reed, D.W., Faulkner, D.S., and Stewart, E.J. 1999. The FORGEX method of rainfall growth estimation. II: Description. *Hydrology and Earth System Sciences*. **3**: 197-203.
30. Robson, A. and Reed, D. 1999. *Flood Estimation Handbook. Volume 3. Statistical procedures for flood frequency estimation*. Institute of Hydrology, Wallingford, 338pp.
31. Weather Log, October 1993. Royal Meteorological Society publication.
32. Weather Log, September 1995. Royal Meteorological Society publication.
33. Wigley, T.M.L., Lough, J.M. and Jones, P.D. 1984. Spatial patterns of precipitation in England and Wales and a revised, homogenous England and Wales precipitation series. *Journal of Climatology*. **4**: 1-25.
34. Zhai, P.M., Sun, A.J., Ren, F.M., Liu, X.N., Gao, B. and Zhang, Q. 1999. Chances of climate extremes in China. *Climatic Change*. **42**: 203-218.

List of Figures

Figure 1. Location of the 204 UK daily rainfall records with complete or almost complete data for the 1961-2000 period and the nine coherent rainfall regions. The regions are: North Scotland (NS), East Scotland (ES), South Scotland (SS), Northern Ireland (NI), Northwest England (NWE), Northeast England (NEE), Central and Eastern England (CEE), Southeast England (SEE) and Southwest England (SWE).

Figure 2. Comparison of mean L-CV and L-Skewness in the nine regions.

Figure 3. Shape of standardized annual maximum distributions for the nine pooling regions.

Figure 4. Fitted standardized annual maximum GEV distributions for the 9 regions.

Figure 5. Fitted annual maximum GEV distributions for the nine pooling regions (using mean regional RMED from 1961-1990 as a multiplier).

Figure 6. Changing L-moment ratios from 1961-2000 for; (a) SWE region, 1-day; (b) NI region, 1-day; (c) CEE region, 10-day; (d) ES region, 10-day.

Figure 7. Decadal change in the 10-day growth curve: (a) SEE; (b) SWE; (c) NEE; (d) NWE.

Figure 8. Decadal change in the 1-, 2-, 5- and 10-day growth curves in the ES pooling region.

Figure 9. Decadal change in RMED: (a) SWE 1-day; (b) NI 1-day; (c) SEE 1-day; (d) SEE 2-day.

Figure 10. Decadal changes in return period estimates using mean regional RMED for the ES pooling region.

Figure 11. Decadal changes in 10-day duration return period estimates using mean regional RMED for: (a) NS; (b) SS; (c) NEE; (d) NWE.




Figure 1. Location of the 204 UK daily rainfall records with complete or almost complete data for the 1961-2000 period and the nine coherent rainfall regions. The regions are: North Scotland (NS), East Scotland (ES), South Scotland (SS), Northern Ireland (NI), Northwest England (NWE), Northeast England (NEE), Central and Eastern England (CEE), Southeast England (SEE) and Southwest England (SWE).




Figure 2. Comparison of mean L-CV and L-Skewness in the nine regions.




Figure 3. Shape of standardized annual maximum distributions for the nine pooling regions.




Figure 4. Fitted standardized annual maximum GEV distributions for the 9 regions.




Figure 5. Fitted annual maximum GEV distributions for the nine pooling regions (using mean regional RMED from 1961-1990 as a multiplier).




Figure 6. Changing L-moment ratios from 1961-2000 for; (a) SWE region, 1-day; (b) NI region, 1-day; (c) CEE region, 10-day; (d) ES region, 10-day.




Figure 7. Decadal change in the 10-day growth curve: (a) SEE; (b) SWE; (c) NEE; (d) NWE.




Figure 8. Decadal change in the 1-, 2-, 5- and 10-day growth curves in the ES pooling region.




Figure 9. Decadal change in RMED: (a) SWE 1-day; (b) NI 1-day; (c) SEE 1-day; (d) SEE 2-day.




Figure 10. Decadal changes in return period estimates using mean regional RMED for the ES pooling region.




Figure 11. Decadal changes in 10-day duration return period estimates using mean regional RMED for: (a) NS; (b) SS; (c) NEE; (d) NWE.

Table 1. Fitted GEV parameters and L-moment ratios for the nine pooling regions using data from 1961-2000.

Pooling Region	Duration	GEV parameters			L-moment ratios		
		ξ	α	k	L-CV	L-skewness	L-kurtosis
SWE	1-day	0.921	0.231	-0.121	0.166	0.250	0.178
	2-day	0.933	0.216	-0.096	0.153	0.233	0.181
	5-day	0.937	0.191	0.011	0.125	0.163	0.167
	10-day	0.951	0.185	0.068	0.116	0.127	0.143
SEE	1-day	0.897	0.247	-0.147	0.185	0.268	0.207
	2-day	0.912	0.224	-0.169	0.172	0.283	0.226
	5-day	0.905	0.21	-0.063	0.149	0.211	0.185
	10-day	0.924	0.22	-0.023	0.148	0.185	0.166
CEE	1-day	0.918	0.27	-0.085	0.186	0.226	0.169
	2-day	0.932	0.252	-0.085	0.172	0.226	0.160
	5-day	0.943	0.243	-0.022	0.158	0.184	0.130
	10-day	0.932	0.222	0.038	0.142	0.146	0.128
NWE	1-day	0.899	0.218	-0.103	0.160	0.238	0.183
	2-day	0.923	0.213	-0.079	0.150	0.222	0.170
	5-day	0.933	0.185	-0.055	0.129	0.206	0.190
	10-day	0.935	0.172	0.012	0.114	0.162	0.188
NEE	1-day	0.897	0.27	-0.064	0.186	0.212	0.155
	2-day	0.915	0.296	-0.014	0.190	0.179	0.138
	5-day	0.925	0.251	-0.036	0.167	0.193	0.155
	10-day	0.931	0.223	-0.043	0.150	0.198	0.140
NI	1-day	0.905	0.228	-0.154	0.173	0.273	0.193
	2-day	0.929	0.245	-0.085	0.170	0.226	0.163
	5-day	0.947	0.207	-0.033	0.138	0.191	0.129
	10-day	0.965	0.171	-0.006	0.112	0.174	0.147
NS	1-day	0.927	0.23	-0.023	0.154	0.185	0.158
	2-day	0.917	0.221	0.008	0.146	0.165	0.166
	5-day	0.949	0.204	0.034	0.129	0.148	0.168
	10-day	0.955	0.193	0.057	0.120	0.134	0.151
SS	1-day	0.943	0.199	-0.036	0.134	0.193	0.162
	2-day	0.951	0.188	-0.037	0.127	0.194	0.193
	5-day	0.962	0.184	-0.006	0.120	0.174	0.168
	10-day	0.963	0.183	0.092	0.111	0.112	0.143
ES	1-day	0.937	0.256	-0.054	0.171	0.205	0.175
	2-day	0.94	0.255	-0.037	0.156	0.179	0.137
	5-day	0.938	0.246	-0.054	0.164	0.205	0.156
	10-day	0.94	0.235	-0.045	0.157	0.199	0.176

Table 2. PUM values for each rainfall duration and each of the nine regions. Bold italicised values give an estimate of the regional PUM in millimetres for the return period (i.e. PUM multiplied by regional mean RMED).

Duration	Return Period (years)	SWE	SEE	CEE	NWE	NEE	NI	NS	SS	ES									
		<i>PUM (mm)</i>	<i>PUM (mm)</i>	<i>PUM (mm)</i>	<i>PUM (mm)</i>	<i>PUM (mm)</i>	<i>PUM (mm)</i>	<i>PUM (mm)</i>	<i>PUM (mm)</i>	<i>PUM (mm)</i>									
1-day	2	0.04	1.5	0.03	0.8	0.03	0.9	0.05	1.8	0.04	1.4	0.05	1.7	0.04	1.6	0.04	1.6	0.04	1.5
	5	0.06	2.2	0.05	1.6	0.05	1.5	0.07	2.7	0.05	1.7	0.07	2.5	0.06	2.3	0.05	2.2	0.05	1.9
	10	0.08	3.1	0.08	2.6	0.07	2.3	0.09	3.7	0.06	1.9	0.09	3.2	0.08	3.2	0.07	3.0	0.08	2.8
	25	0.13	4.9	0.13	4.4	0.11	3.5	0.13	5.4	0.07	2.3	0.12	4.3	0.11	4.7	0.10	4.4	0.12	4.1
	50	0.18	6.6	0.19	6.1	0.15	4.5	0.17	7.0	0.09	2.9	0.15	5.5	0.14	6.0	0.13	5.7	0.16	5.5
2-day	2	0.04	1.8	0.04	1.5	0.04	1.5	0.04	2.1	0.05	2.0	0.04	2.0	0.04	2.6	0.03	1.6	0.05	2.1
	5	0.06	2.6	0.06	2.4	0.07	2.7	0.06	3.2	0.06	2.6	0.07	3.4	0.07	4.0	0.04	2.2	0.06	2.9
	10	0.07	3.4	0.09	3.8	0.10	3.7	0.09	4.5	0.08	3.2	0.10	4.9	0.09	5.5	0.06	3.3	0.08	3.8
	25	0.10	4.7	0.16	6.4	0.14	5.3	0.13	6.9	0.11	4.5	0.15	7.2	0.13	8.0	0.10	5.4	0.12	5.6
	50	0.13	6.0	0.22	8.9	0.17	6.7	0.17	9.1	0.14	5.6	0.20	9.3	0.17	10.0	0.13	7.4	0.14	6.8
5-day	2	0.03	2.4	0.03	1.9	0.03	1.5	0.03	2.4	0.04	2.0	0.03	2.1	0.04	3.8	0.03	2.5	0.04	2.5
	5	0.05	3.7	0.04	2.5	0.05	2.5	0.05	4.1	0.06	3.2	0.06	3.8	0.07	6.3	0.04	3.2	0.05	3.5
	10	0.08	5.3	0.07	4.2	0.07	3.6	0.08	6.0	0.08	4.3	0.08	5.6	0.09	8.4	0.05	4.3	0.08	5.4
	25	0.11	7.9	0.12	7.0	0.10	5.2	0.12	9.3	0.11	6.0	0.12	8.2	0.13	12.2	0.08	6.5	0.13	8.8
	50	0.14	10.0	0.16	9.5	0.13	6.7	0.16	12.3	0.13	7.5	0.16	10.5	0.16	14.9	0.10	8.7	0.17	11.3
10-day	2	0.04	3.8	0.04	3.0	0.03	2.1	0.02	2.5	0.04	2.9	0.03	2.4	0.04	4.8	0.02	2.2	0.03	2.9
	5	0.05	5.2	0.04	3.4	0.05	3.5	0.03	3.7	0.06	4.8	0.05	4.3	0.06	8.1	0.03	4.4	0.05	4.8
	10	0.07	6.6	0.06	4.8	0.07	4.8	0.05	5.5	0.08	6.1	0.07	6.2	0.08	11.1	0.05	6.2	0.08	7.1
	25	0.10	9.7	0.09	7.4	0.10	6.8	0.08	9.0	0.10	7.8	0.10	8.9	0.12	15.8	0.07	9.2	0.13	11.3
	50	0.12	12.5	0.13	10.2	0.12	8.4	0.11	12.5	0.12	9.3	0.13	11.5	0.14	19.2	0.09	11.5	0.16	14.1

Table 3. Mean RMED per decade for the nine pooling regions for each of: (a) 1-day; (b) 2-day; (c) 5-day; and (d) 10-day; durations. Highest value of decadal RMED shown in bold italic.

Pooling Region	1961-70	1971-80	1981-90	1991-00
(a)				
SWE	41.4	40.3	37.7	39.1
SEE	36.9	37.2	32.2	33.5
CEE	34.3	32.8	32.1	34.6
NWE	42.8	43.4	42.3	41.0
NEE	34.5	34.2	36.8	34.3
NI	42.4	36.2	38.2	36.6
NS	44.1	42.2	47.2	45.1
SS	44.1	42.9	44.3	47.7
ES	37.6	33.8	40.5	42.1
(b)				
SWE	51.9	53.0	49.2	52.0
SEE	48.6	45.5	41.4	41.1
CEE	41.9	40.9	41.5	44.1
NWE	57.0	57.6	55.1	55.3
NEE	42.9	44.1	46.9	45.7
NI	54.2	48.8	51.6	51.0
NS	61.5	58.7	65.6	63.8
SS	58.2	56.7	60.0	63.9
ES	49.8	47.3	50.6	56.7
(c)				
SWE	71.2	74.4	75.1	76.6
SEE	62.1	62.2	58.8	58.0
CEE	52.6	54.6	53.7	57.3
NWE	82.8	82.9	81.4	86.2
NEE	56.7	59.9	60.9	62.9
NI	73.9	68.5	69.8	75.8
NS	90.9	91.0	105.3	102.4
SS	87.6	83.2	95.4	101.4
ES	67.9	63.6	70.7	82.7
(d)				
SWE	101.2	103.1	107.6	108.2
SEE	82.5	84.5	85.0	82.0
CEE	70.2	73.8	71.8	74.3
NWE	118.5	115.8	116.4	120.8
NEE	76.4	82.1	80.1	83.0
NI	96.1	94.0	97.4	103.0
NS	130.6	134.9	151.7	152.2
SS	126.8	122.4	141.6	147.0
ES	91.9	86.3	96.8	110.5

Table 4. Change in the 100-year rainfall event for: (a) 5-day; and (b) 10-day; durations for 1961-1990 and 1991-2000. Italic indicates lower extremes in 1991-2000 than previously.

Pooling Region	Rainfall from the 100-year Event (mm) (1961-1990)	Rainfall from the 100-year Event (mm) (1991-2000)	% Change
(a)			
SWE	128	129	0
<i>SEE</i>	<i>123</i>	<i>104</i>	<i>-15</i>
CEE	105	106	2
NWE	148	160	9
NEE	115	130	13
NI	131	143	9
NS	159	171	8
SS	144	177	23
ES	122	208	71
(b)			
<i>SWE</i>	<i>175</i>	<i>160</i>	<i>-8</i>
<i>SEE</i>	<i>163</i>	<i>133</i>	<i>-19</i>
<i>CEE</i>	<i>127</i>	<i>123</i>	<i>-3</i>
NWE	185	206	11
NEE	143	168	18
<i>NI</i>	<i>161</i>	<i>158</i>	<i>-2</i>
NS	219	233	6
SS	196	224	15
ES	159	274	72

Table 5. Probability of occurrence of the 1991-2000: (a) 5-day; (b) 10-day; rainfall event corresponding to the 1961-1990 50-year event.

Pooling Region	Probability of event (%)	Return period
(a)		
SWE	2.2	45
SEE	0.6	167
CEE	2.9	35
NWE	3.4	30
NEE	3.9	26
NI	3.6	28
NS	3.5	28
SS	8.6	12
ES	13.5	7
(b)		
SWE	0.8	130
SEE	0.2	500
CEE	1.9	53
NWE	4.2	24
NEE	4.9	21
NI	2.4	42
NS	4.5	22
SS	9.2	11
ES	12.9	8